
The effect of plant-density and sowing-date on yield of Basil (*Ocimum basilicum* L.) In Iran

Sedigheh Sadeghi^{1*}, Aptin Rahnavard² and Zoheir Y. Ashrafi^{1,3}

¹Department Weed Science, Faculty of Agronomy and Animal Science, University College of Agricultural and Natural Resource, Karaj Paradise, University of Tehran, Iran.

²Islamic Azad University, Tonekabon branch, Iran.

³Payame Noor University, Tehran State, Iran.

Sadeghi, S., Rahnavard, A. and Ashrafi, Z.Y. (2009). The effect of plant-density and sowing-date on yield of Basil (*Ocimum basilicum* L.) in Iran. Journal of Agricultural Technology 5(2): 413-422.

The experiment was conducted to evaluate the effect of sowing date and plant density on yield and yield components of Basil (*Ocimum basilicum* L.) under dry farming conditions. Four plant densities (50, 100, 150 and 200 plants m⁻²) and three sowing dates (3, 13 and 23 of March) were applied. Result showed that seed yield was influenced by sowing date and plant density interaction. Early sowing date resulted in higher seed yields as evident from higher aboveground biomass, the number of umbrella per plant, the number of seed per umbrella and plant height. Harvest index and 1000-seed weight was not affected by sowing date and planting density. Earlier sown plants with density of 200 m⁻² resulted in higher seed yields.

Key words: Sowing date, plant density, basil, yield, yield components

Introduction

Basil (*Ocimum basilicum* L.) is a member of Apiaceae. This species originated in Egypt and East Mediterranean, but is widely cultivated in Iran, Japan, China and Turkey. Basil has a long history of use as food flavors, perfumes and medicinal values. Essential oil has been used for bringing smell to some medicines, for sterilizing of surgical operation fiber, for producing of some veterinary and agricultural medicines and plastic (Simon *et al.*, 1984). Basil seeds have an aromatic odor and bitter taste. They are used as an essential ingredient in soup, sausages, cheese, cakes and candies (Behera *et al.*, 2004). Presently, Iran is an important Basil exporter, constituting 20-40% of world market (Barros *et al.*, 2004). In semiarid area such as Iran, water is the most limiting factor for farming. Basil can be done some as fallow crop in wheat or

*Corresponding author: Sedigheh Sadeghi; e-mail: sadeghi58@gmail.com

barley fallow in dry land farming of Iran. In suitable plant density, plants are completely use in environmental conditions (water, air, light and soil) and inter- or intra-specific competition is minimum. Yield loss due to unfavorable sowing date has been reported in many crops such as sunflower (Barros *et al.*, 2004) and fennel (Bianco *et al.*, 1994; Kafi, 1990). Yield components of Basil include the number of plant in area unit, the number of umbrella per plant, the number of seed per umbrella and seed weight. The number of plant per unit area is the most important among yield components (Kafi, 2003). Ahmed and Haque (1986) studied the effect of row spacing (15, 20, 25 and 30 cm) and time of sowing (November 1, November 20, December 10 and December 30) on the yield of black cumin (*Nigella sativa*) in Bangladesh, they found that closer row spacing (15 cm) and early sowing (November 1) was the best for higher seed yield of black cumin. The number of umbrella per plant has the second rank interms of importance in yield components. Aminpour and Karimi (1995) reported that 96% of seed yield variation was related to this yield component. The number of seed per umbrella is affected by environmental, field management and its number was reported from 11.3 to 16.8 under varying plant densities (Kafi, 2003). The weight of Basil seed varied in different experiments. Kafi (2003) reported that it was from 2.79 to 2.99 g under varying plant densities. Shortening of the growing cycle decreased the amount of radiation intercepted during the growing season and thus total dry weight of plant (Andrade, 1995). With delayed sowing, development is accelerated because the crops encounter higher temperatures during the vegetative growth (Damato *et al.*, 1994). Ehteramian (2003) reported that delayed sowing date was better because of the occurrence, resulting to lack sudden winter chilling. Delayed sowing date decreases seed weight and the number of umbrella per plant. Optimum plant density and sowing date of Basil in Khorasan province of Iran has not yet been investigated. This study aimed to determine the determination of the optimum sowing date and plant density of Basil for achievement of maximum seed yields under the conditions of the west of Iran.

Materials and methods

The experiment was conducted in 2005 at Research Farm, Faculty of Agriculture, Tehran University, Karaj, Iran (latitude 35° 50' N, longitude 50° 58'E, and altitude 132 m). Long term average precipitation was 456 mm and this area is semiarid according to De Martonne classification (Ref). In the experiment area clay soil texture was used with soil pH of 7.64 and EC 1.3 dS m⁻¹.

Experiment design and cultural practices

In this experiment Randomized Complete Block Design was used and divided into four replications. Three sowing dates (3, 13 and 23 of March) was applied with four plant densities (50, 100, 150 and 200 plants m⁻²), respectively. Row spacing was measured at 0.3 m and with the plot length of 2.1 m, plot wideness 2.1 m and plot area 4.41 m². The distances between plots and between blocks was 1 m and 2 m respectively. According to meteorological statistics of the area, there was no danger of chilled at this time. Khorasan cultivar was used. The seedling depth 0.5 cm before sowing and the soil was leveled at 100 kg ha⁻¹ of diamonium phosphate (DAP) as a fertilizer was applied. Planting of seeds was done manually. At 3-4 leaf stage, the seedlings were thinned with expected densities. Hand weeding was applied for irradiation of weeds and insect control with the used of Sevin insecticide.

Plant identification and statistical analysis

In order to determine the yield and other characters under varying plant densities and sowing dates, a number of plant samples was taken by 1 m² quadrat (Latond, 1994). Up to 50 cm, primer and edge lines were discarded and four planting rows were harvested. All plots were evaluated on 1 m² area. In order to measure the seed yield and total dry matter, plants was cut and after drying, dry matter and seed yield was measured. Plants were harvested at physiological maturity stage when plants turn into yellowed. Six plants was randomly were selected in each plot to measure the number of umbrella per plant and plant height. Three umbrellas in each plant (from six selected plants as mentioned above) were selected and the number of seed was calculated. Harvest index was computed as the ratio of the seed yield to aboveground dry matter at harvest. Analysis of variance (ANOVA) was used to determine the significant differences. The Multiple Range Test of Duncan performed the separation of means when the *F* test revealed the error probability to justify the difference minor. Correlation coefficients were calculated for the relationship between seed yield and several crop parameters. All statistics was performed with MSTATC program (version 2.10) and SPSS (version 10.0).

Results and discussion

Seed yield

The effects of sowing date, plant density and their interaction on seed yield were significant (Table 1). The first sowing date, plants with density of 200 plants m⁻² had the highest seed yield. However, the yield at 150 plants m⁻²

density was no significant difference compared to 200 plant m⁻² density. Under the second sowing date (D2), studied that the plant densities except plant densities of 150 and 200 plants m⁻² have no significant differences. Under the third sowing date (D3), plant density of 150 plants m⁻² resulted in higher seed yield; therefore, no significant difference from plant density of 200 plants m⁻². Plant densities of 50 and 100 plants m⁻² had a yield reduction of 58% and 42% respectively compared with 150 plants m⁻² (Table 2). Plant density of 200 plants m⁻² under D1 and D2 resulted in higher seed yields however, the plant density have no significant difference from 150 plants m⁻² under D1 and 50 and 100 plants m⁻² under D2 . Under D3, plant densities of 150 to 200 plants m⁻² had the highest seed yields. El-Gengai and Abdallah (1978) and Bianco *et al.* (1994) reported significant effect of sowing date and plant density on seed yield of fennel (*Foeniculum vulgare* Mill.). Results obtained from the study were comparable with Ehteramian (2003) findings, but contrary to those of Damato *et al.* (1994). There was a positive correlation between seed yield and aboveground biomass($r=0.91^{**}$). Early sowing dates resulted in higher seed yields that can be explained by higher aboveground biomass, the number of umbrella per plant, the number of seed per umbrella and plant height (Table 2). In view of the sensitivity of Basil to climatic factors especially to photoperiod and temperature, it is essential that sowing should be done on time so that there have enough time for vegetative growth. Delayed sowing results in reduced vegetative growth leading to reduced number of umbrella per plant and plant height (Okut, 2001; Rahimian, 1991; Sharratt and Gesch, 2004). Under optimum plant density, plants show efficient use of available water, light and nutrient while under high plant density, and have a higher competition among plants.

Table 1. Analysis of variance results (Mean of Square) for different traits of basil under varying sowing dates and plant densities.


Trait	sowing date	plant density	plant density * sowing date
Seed yield	429.165**	216.894**	153.101**
Aboveground biomass	4172.934**	1839.311**	611.679**
Number of Leaf per plant	109.841*	47.079	36.385
Number of seed per umbrella	3.468**	2.643**	2.576**
1000-seed weight	00.201	0.082	0.046
Harvest index	0.156	0.051	0.065
Plant height	20.976**	0.504	4.726*

* Significant at the 0.05 level.

** Significant at the 0.01 level.

Aboveground biomass

The effect of sowing date, plant density and their interaction were significant for aboveground biomass (Table 1). The aboveground biomass showed an increasing trend, with increases in plant density under the first sowing date (D1), plant density of 200 plants m⁻² had the highest aboveground biomass (Table 2). There was strongly correlation ($r=0.91^{**}$) between seed yield and aboveground biomass, but a negative one ($r=-0.68^*$) between aboveground biomass and harvest index. The effect of sowing date and plant density interaction on aboveground biomass was favourable to high seed yield.



#sowing dates of 1, 2, and 3 are 3, 13 and 23 of March respectively.

Fig. 1. Mean comparisons for the number of leaf per plant under different sowing dates Means followed by the same letter are not significantly different at $P=0.05$ (*) or $P=0.01$ (**) according to Duncan Multiple Range Test.

Yield components

The effect of plant density on the number of umbrellas per plant had no significant (Table 1), due to compensatory capacity of the other yield components such as the number of seed per umbrella. With changing plant density, each plant changes the number of seed per umbrella that results in fixed number of umbrellas per plant. There has no significant correlation between the number of umbrellas per plant and seed yield ($r=0.16$ ns). Ehtramian (2003) reported that correlation between seed yield and the number of umbrellas per plant was $r=0.22$. Bianco *et al.* (1994) found significant effect of plant density on the number of umbrellas per plant. In this research, average number of umbrellas per plant was 16.56. Kafi (2003) reported that the number of umbrellas per plant under varying plant densities was from 18.9 to 31.3. In this study the present results was lower than those of Tuncturk and Tuncturk (2006), Arslan and Bayrak (1987) and Okut (2001). These differences was probably do to the variations in environmental conditions, genotype and soil

properties. Sowing date had the significant effect on the number of umbrella per plant (Table 1) that related to high sensitivity of this yield component to photoperiod and temperature, which conforms to the finding of Rahimi (1993). The highest number of umbrella per plant was achieved under the first sowing date (give actual sowing dates). Three sowing dates were 3, 13 and 23 on March. Plants under second (13, March) and third (23, March) sowing dates had 25 and 18% lower umbrella number compared with the first sowing date (3, March) (Fig. 1). The effect of sowing date and plant density and their interaction on the number of seed per umbrella was significantly' statistically (Table 1). The number of seed per umbrella showed a decreasing trend, with decreases in plant densities and the plant density of 50 plants m^{-2} had the highest number of seed per umbrella and the plant densities of 100 and 150 plants m^{-2} had no significant difference for the number of seed per umbrella (Table 2). Because seed set depends on sufficient nutrients they provide and environmental conditions while shift from vegetative to reproductive stage, increased plant densities result in limited availability of nutrients, light and water so the number of reproductive units decrease; at last seed number reproduction decreases. Kafi (2003) found that the number of seed per umbrella under different plant densities (40, 80 and 200 plants m^{-2}) decreased. That results was comparable to the finding of Kafi (2003), who noted that no correlation between the number of seed per umbrella and seed yield ($r=-0.006$). Ehteramian (2003) reported that correlation between seed yield and the number of seed per umbrella was ($r=0.75$). The first sowing date (D1) except under plant densities of 50 and 200 plants m^{-2} had got the highest number of seed per umbrella in relation to D2 and D3. It was due to sensitivity of the number of seed per umbrella to photoperiod and temperature (Table 2). Environmental conditions during pollination at the first stage of seed set, determine the number of seed per umbrella (16 quote properly). The effect of sowing date and plant density on 1000-seed weight was found not significant (Table 1.), which seemed to be more dependent on genetic than environmental factors (18 quote properly). These data were in contrast results of Barros *et al.* (2004) and Tuncturk and Tuncturk (2006). Data indicated no correlation between seed yield and 1000-seed weight ($r=0.429$), but a negative one between 1000- seed weight and plant height ($r=-0.677^*$). In this study, 1000-seed weight was 2-3 g, which was lower than that reported by Arslan and Bayrak (1987), Okut (2001) and Tuncturk and Tuncturk (2006), but similar to that of Kafi (2003).

Harvest index and plant height

The effect of sowing date and plant density on harvest index has no significant (Table 1), to change plant density or sowing date. Changes in reproductive and vegetative parts had the same rate, as changed the plant density or sowing date. The decrease or increase in aboveground biomass of single plant was consistent with changes in seed yield per plant (Table 2). Ball *et al.* (2000), Behera *et al.* (2004) and Kafi (2003) also reported the same result. There was no correlation between harvest index and seed yield, but on this harvest index with aboveground biomass was negative ($r=-0.608$). The average harvest index of this study was 0.45 that different from the report of Ehteramian (2003). The effect of sowing date, plant density and their interaction on plant height was significant (Table 1) and delayed sowing decreased plant height (Table 2). The first sowing date (D1 give actual date) except under plant density of 50 and 100 plants m^{-2} resulted in higher plant height than third sowing date (D3 give actual date). It might be explained by a higher dry matter accumulation and vegetative growth due to early sowing, while delayed sowing reduced plant height due to high sensitivity to photoperiod and temperature. Under early sowing date, plant densities of 50, 100 and 150 plants m^{-2} had a plant height reduction of 16, 10 and 5% compared with plant density of 200 plants m^{-2} . Such an increase in plant height with increased plant density may be explained by increased activity of stem growth hormone due to light deficiency (10 quote properly). El-Gengai and Abdallah (1978) and Bianco *et al.* (1994) found that sowing date and plant density had significant effect on plant height of fennel. There was no correlation between seed yield and plant height ($r=0.455$), while Correlation of plant height and 1000-seed weight was negative ($r=-0.677^*$).

Conclusions

Basil is sensitive to plant density and sowing date. Early sowing in dry land Basil was results in critical to increase in seed yield, possibly due to higher aboveground biomass, the number of umbrella per plant and plant height. Lower densities do not produce sufficient seed per unit area. However, the relatively small absolute differences in seed yield between some plant densities demonstrated the remarkable compensation capacity of Basil between the different yield components.

Acknowledgment

The authors express their appreciation to University of Tehran and Azad Islamic University for full support of the project.

References

- Ahmed, N.U. and Haque, K.R. (1986). Effect of row spacing and time of sowing on the yield of black cumin (*Nigella sativa* L.). Bangladesh. J. Agric. 1: 21-4.
- Aminpour, R., and Karimi, M. (1995). Effect of irrigation number on water potential, yield and yield components of Basil (*Basilum cyminum*). Master of Science Thesis of Agronomy, Isf. Univ. Technol., Isfahan, Iran.
- Andrade, F.H. (1995). Analisis of growth and yield of maize, soybean grown at Balcare, Argentina. Field Crop Res. 41: 1-12.
- Arslan, N. and Bayrak, A. (1987). Effects on yield and some yield components of Basil (*Ocimum basilicum*L.) of different sowing times. Doğa Tu Agr.And For. Jour. 11, 2, Ankara, Turkey.
- Ball, R.A., Purcell, L.C. and Vories, E.D. (2000). Short-season soybean production and yield compensation in response to plant population and water regime. Crop Science 40: 1070-1078.
- Barros, J.F.C., Carvalho, M.D. and Basch, G. (2004). Response of sunflower (*Helianthus annus* L.) to sowing date and plant density under Mediterranean conditions. Europ. J.Agronomy. 21: 347-356.
- Behera, S., Nagarajan, S. and Rao, L.J.M. (2004). Microwave heating and conventional roasting of Basil Seeds (*Ocimum basilicum*L.) and effect on chemical composition of volatiles. Food Chemistry 87(1): 25-29.
- Bianco, V.V., Damato, G. and Gridi, A. (1994). Sowing date, plant density and crowing cutting on yield and quality of Florence fennel seed. International Symposium on Agrotechnics and Storage of Vegetative and Ornamental Seeds. Bari, Italy.
- Damato, G., Biaco, V.A and Laterza, M. (1994). First result of plant density and nitrogen rate on yield and quality of Florence fennel (*Foeniculum vulgar* Mill. Var. Azotericum Thell.) seeds. International Symposium on Agrotechnics and Storage of Vegetative and Ornamental Seeds. Bari, Italy. July.
- Ehteramian, K. (2003). The effects of different levels of nitrogen fertilizer and plant dating on Basil (*Ocimum basilicum*L.) in Kooshkak region in the Fars province. Master of Science Thesis of arid area management. Shiraz Univ., Shiraz., Iran.
- El-Gengai, S and Abdallah, N. (1978). The effect of date of sowing and plant spacing on yield of seed and volatile oil of fennel (*Foeniculum vulgar* Mill.). Pharmazie 9: 605- 6.
- Kafi, M. (1990). Study on weed control number, row spacing and plant density on growth and yield of Basil (*Ocimum basilicum*L.). Master of Science Thesis of Agronomy, Ferdowsi Univ. Mashhad., Mashhad, Iran.
- Kafi, M. (2003). Basil- Production and Processing . Ferdowsi Univ. Mashhad Publication. Mashhad, Iran. 195 pp.
- Latond, G.P. (1994). Effect of row spacing, seeding rate and nitrogen on yield of barley and wheat under zero-till management. Canadian Journal of Plant Science 74: 703-711.
- Mollafiani, A.A. (1992). Effect of sowing date and row spacing on yield of cumin under dry land and rainfed farming. Iranian Research Organization for Science and Technology. Khorasan Reseach Center. Khorasan, Iran.
- Okut, N. (2001). Effects of different nitrogen doses and row spaces on the yield and quality components of cumin (*Cuminum cyminum* L.).PhD Thesis. Yüzüncü Yıl Univ. Van, Turkey.
- Rahimi, M. (1993). Study on possibility of chemical weed control of Basil. Iranian Research Organization for Science and Technology. Khorasan Reseach Center. Khorasan., Iran.

- Rahimian, M.H. 1991. Effect of sowing date and irrigation regime on growth and yield of Basil. Iranian Research Organization for Science and Technology. Khorasan Research Center. Khorasan., Iran.
- Sharratt, B.S. and Gesch, R.W. (2004). Water Use and Root Length Density of *Cuphea* spp. Influenced by Row Spacing and Sowing Date. *Agron. J.* 96: 1475-1480.
- Simon, J.E., Chadwick, A.F. and Crack, L.E. (1984). *Herbs: An Indexed Bibliography. 1971-1980. The scientific literature on selected herbs, and aromatic and medicinal plants of the temperate zone.* Arcon Book press, Hamden CT, 770 PP
- Tunçturk, R. and Tunçturk, M. (2006). Effects of Different Phosphorus Levels on the Yield and Quality Components of Cumin (*Cuminum cyminum* L.). *Res. J. Agric. & Biol. Sci.* 2(6): 336-340.

(Received 9 September 2008; accepted 20 July 2009)

Table 2. Mean comparisons for different traits of cumin under varying sowing dates and plant densities.

Plant density	Seed yield (g m ⁻²) **			Plant height (cm)*			Aboveground biomass (g m ⁻²) **			number of seed per umbrella**		
	Sowing date [#]			Sowing date			Sowing date [#]			Sowing date		
	1	2	3	1	2	3	1	2	3	1	2	3
50	24.96 cd	27.63abc	11.47e	12.62bc	12.99 abs	12.27bc	61.95bc	67.00bc	21.9e	6.930ab	7.475a	6.740abc
100	21.46 cde	25.61 bcd	16.07de	13.46 abc	12.27 bc	12.36bc	63.75bc	47.81cd	25.71de	6.922ab	5.675cde	5.355e
150	27.73 abc	19.52cde	27.58abc	14.19ab	13.4abc	11.59cd	62.55bc	56.17bc	60.78bc	7.280a	5.185e	5.995bcde
200	36.64 a	35.36ab	18.53cde	15.01a	12.95abc	9.97d	93.7a	78.55ab	49.04cd	6.485abcd	5.570de	7.415a

Means followed by the same letter are not significantly different at $P=0.05$ (*) or $P=0.01$ () according to Duncan Multiple Range Test.

[#]Sowing dates of 1, 2, 3 are 3, 13 and 23 of March respectively.