
Activity Level of Lactate Dehydrogenase and B-Glucosidase Enzymes in the Honeybee Colonies, (*Apis Mellifera* L.) with Different Feeding

Mostafa, S. *, El Naby, A. and Zidan, E. W.

Plant Protection Research Institute (PPRI), ARC, Dokki, Giza.

Mostafa, S., El Naby, A. and Zidan, E. W. (2014). Activity level of lactate dehydrogenase and β -glucosidase enzymes in the honeybee colonies, (*Apis mellifera* L.) with different feeding. International Journal of Agricultural Technology 10(2):483-491.

Abstract The activity of Lactate dehydrogenase and β -glucosidase was determined in Carniolan workers honey bee, *Apis mellifera* L. after feeding on different diets antibiotic ,ripe milk and some natural plant extract (thyme, clove, cinnamon, black seeds and anise) . The results showed the highest increase in the β -glucosidase activity was occurred in the diet containing Ripe milk (106.17%) and the other diets scored decrease in the activity , as well as the Lactate dehydrogenase activity the results showed highly enzyme activity in diets antibiotic and clove 192.56%, & 215.70% respectively as compared with the other diets used in this study.

Keywords: Honey bee, β -glucosidase, Lactate dehydrogenase, antibiotic, natural plant extract and ripe milk.

Introduction

Carbohydrates and fatty acids play a predominant role in the generation of energy for prolonged flights in insects (Beenackers, 1969; Sacktor, 1970). Insects utilize various high-energy fuels for flight. They derive energy to meet the intense metabolic needs of flight from carbohydrates, fat or amino acids such as proline (Wheeler, 1989; Candy, 1989). During flight, honeybees increase their metabolic rate to relatively high values (Harrison and Fewell 2002; Feuerbacher *et al.*, 2003). In honey bee colonies requires proteins, carbohydrates, lipids, vitamins, minerals and water. These nutrients must be in the diet in a definite qualitative the food metabolism plays an important role in bee life. The physiological digestion different in bee body composition; percentage of glycogen, lipids and proteins (Hrassing and Crailsheim, 2005). The source of feeding honey bees effects on induction haemolymph enzymes (Bounias and Morgan, 1990).

* **Crossbonding author:** El-Mohamedy, R. S. R; **Email:** riadelmohamedy@yahoo.com

The enzyme glucosidase is ubiquitous and occurs in all the living kingdoms starting from bacteria to highly evolved mammals and performs varied functions in these organisms. In bacteria and fungi, β -glucosidases are mainly a part of the cellulase enzyme system and are responsible for the hydrolysis of short chain oligosaccharides and cellobiose (Bisaria and Mishra, 1989; Kubicek *et al.*, 1993). In insects and plants, β -glucosidase is involved in the release of cyanides from cyano-glucoside precursors. This is a part of a defense mechanism displayed in these systems (Esen, 1993). Biochemical characteristics of β -glucosidase have been studied in the digestive system and salivary glands of many insect species belong to different orders, such as *Glyphodes pyloalis* Walker (Lep.: Pyralidae), *Apis mellifera* L. (Hymenoptera: Apidae), *Drosophila melanogaster* Meigen (Diptera: Drosophilidae), *Xanthogaleruca luteola* Mull. (Col.: Chrysomelidae) and *Rhynchophorus ferrugineus* Olivieri (Col.: Curculionide) (Ghadamyari *et al.*, 2010; Riseh *et al.*, 2012; Huber and Mathison, 1976; Sharifi *et al.*, 2011; Tanimura *et al.*, 1976; Chipoulet and Chararas, 1985; Pratviel-Sosa *et al.*, 1987; Santos and Terra, 1985; Terra and Ferreira, 1958). β -glucosidase can degrade cellobiose and cello-oligosaccharides to glucose in insects. Hemicelluloses and cellulose present in insect food are converted to di- and oligo- β -saccharides by some carbohydrases and digestive β -glucosidases that play an important role in their hydrolysis (Terra WR, Ferreira, 1994). Also, several investigations indicate the main role of β -glucosidases in insect-host plant interaction (Ferreira *et al.*, 1997; Ferreira *et al.*, 1998; Mattiacci *et al.*, 1995; Marana *et al.*, 2000). β -glucosidases are enzymes that catalyze the cleavage of glycosidic bonds in oligosaccharides or glycoconjugates. Several glucosidases are specific for the cleavage of glycosidic bonds depending on the number, position, or configuration of the hydroxyl groups in the sugar molecule. The activity of glucosidases is fundamental to several biochemical processes such as degradation of diet polysaccharides to furnish monosaccharide units, which are then able to be metabolically and used by the organism, lysosomal glycoconjugate catabolism and glycoprotein processing, and biosynthesis of oligosaccharide units in glycoproteins or glycolipids (Murray *et al.*, 1994).

In insects, the fat body is the main organ responsible for energetic metabolism. This is also the organ of conversion and storage of fat, carbohydrates, and proteins (Arrese and Soulages, 2010). The fat body is responsible for metabolism of carbohydrates and is especially enlarged in insect larvae. In honeybee larvae, it can constitute up to 65% of an individual's body mass (Bishop, 1925).

Lactate dehydrogenase (LDH) is an important glycolytic enzyme present in virtually all animal tissues (Kaplan and Pesce, 1996). It is also involved in

carbohydrate metabolism and has been used to indicate exposure to chemical stress (Diamantino *et al.*, 2001). LDH is involved in the production of energy, being particularly important when a considerable amount of additional energy is required immediately. A negative correlation between LDH activity and ambient oxygen levels for some aquatic organisms were suggesting a possible biochemical adjustment in response to the lowered oxygen levels. This probably occurs also in situations of chemical stress. Therefore, this enzyme may be a sensitive criterion in laboratory (Senthil Nathan *et al.*, 2005). LDH is, also, a parameter widely used in toxicology and in clinical chemistry to diagnose cell, tissue, and organ damage. However, the potential of this enzyme as an indicative criterion in the invertebrate toxicity tests has been scarcely explored (Senthil Nathan *et al.*, 2006).

The aim of this study was to evaluate the effect of different feeding period winter to help bee colonies to surmount on the malnutrition particularly at unsuitable conditions

Materials and methods

Feeding Honeybee Colonies

This work was carried out at the apiary of the Agriculture Research Station, Beni sweif Egypt during the period from 22nd Dec. 2012 to 10th of Feb. 2013. Twenty four healthy honeybee colonies from Carniolian Hybrid were conducted for this study.

Food supplement

A cake of 100gm food supplement consisted of one part medical dried yeast mixed with two parts powder sugar (w/w) according to zidan (2009).

Diet (1) content 100gm of food supplement and 2gm antibiotic (Sodium Sulphate demidin)

Diet (2) content 100gm of food supplement and 30ml of natural plant extract of Thyme.

Diet (3) content 100gm of food supplement and 30ml of natural plant extract of Clove.

Diet (4) content 100gm of food supplement and 30ml of natural plant extract of Cinnamon.

Diet (5) content 100gm of food supplement and 30ml of natural plant extract of black seeds.

Diet (6) content 100gm of food supplement and 30ml of natural plant extract of Anise.

Diet (7) content Ripe milk mixed with powder sugar (1: 3) (w:w).
Diet (8) Control (fed on food supplement without additives).

100 gm of blend was placed on the top of the Combs from three colonies weekly to feed throughout the experimental periods. Samples of 100 adult workers were taken directly from the central comb of each colony at the end experimental. Bee Samples was transferred to the Laboratory and immediately frozen stored until analysis.

Preparation of extracts from bees

The samples were homogenized in a glass Potter homogenizer on an ice bath with 2.5 ml. cool 0.9% NaCl (1:10 w/v). Homogenates were centrifuged for 15 min at 1000 x g at 4 °C. In the supernatants.

Determination of enzymatic activity

Determination of Lactate dehydrogenase activity

The method described here is derived from the formulation recommended by the German Society for clinical chemistry (DGKC, 1972). Lactate dehydrogenase of catalyzes the conversion pyruvate to lactate, NADH is oxidized to NAD in the process. The rate of decrease in NADH is directly proportional to the LDH activity and determined photometrically

Determination of β - glucosidase activity

β - glucosidase activity was measured by assaying glucose liberated by enzymatic hydrolysis of salicin as described by Lindorth (1988).

Results and discussions

Data show in Table 1 indicated increasing percentage in the biological activity to area of sealed worker brood reared, the diets which content antibiotic and ripe milk recorded higher value percentage (429.96% and 384.95, respectively) , the same diets recorded higher value percentage for mean number of combs covered with bees (115.475% and 107.85, respectively) and recorded higher value percentage pollen stored with all diets than control, but found the diets which content antibiotic and ripe milk recorded higher value percentage 38.80% and &33.78%, respectively). Zidan (2009) fed honey bee colonies with food supplement mixed with 6ml of each of the previously prepared mixture; eucalyptus, peppermint and thymol oils by different concentration and the control one was fed only with cake. He found increased

the biological activity the biological activities of treated bee colony (sealed worker brood, covered combs with bee, pollen grain stored, royal jelly and honey), advice the beekeeper to using essential oils with sugar solution added to pollen supplement for their directly that help the bee workers to the reactivation after the season end.

Table 1. Effect of the different feeding on some biological activities of the honey bee Colonies

Treatments	Mean area of sealed worker brood (inch ²)			Mean No. of combs covered with bee /colony			Pollen grain	
	Treatments		Increasing %	Treatments		Increasing %	Increasing %	Mean area (inch ²)
	Before	After		Before	After			
Diet(1)	115.47	9.33	4.33	429.96	1190.67	224.67	276.67	38.80
Diet(2)	100.23	8.67	4.33	328.85	1060.67	247.33	262.67	31.78
Diet(3)	78.37	8.33	4.67	204.66	960.67	315.33	249.33	25.08
Diet(4)	92.38	8.33	4.33	204.24	910.67	299.33	255.33	28.09
Diet(5)	78.37	8.33	4.67	279.70	1035.33	272.67	245.33	23.10
Diet(6)	85.65	8.67	4.67	193.53	966.67	329.33	260.67	30.77
Diet(7)	107.85	9.00	4.33	384.95	1290.67	266.67	266.67	33.78
Diet(8)	64.24	7.67	4.67	138.41	740.67	310.67	199.33	-

Table 2. Enzyme activity for β - glucosidase and Lactate dehydrogenase at homogenate honeybee

Treatment	Lactate dehydrogenase U x 10 ³ /ml			β - glucosidase (ug glucose./min/mg)		
	Mean \pm SE	A*	%	Mean \pm SE	A*	%
Diet (1)	77.667 \pm 0.752	1.93	192.56	2.860 \pm 0.235	0.24	23.95
Diet (2)	32.000 \pm 0.965	0.79	79.34	3.787 \pm 0.133	0.32	31.71
Diet (3)	87.000 \pm 0.769	2.16	215.70	4.153 \pm 0.359	0.35	34.77
Diet (4)	28.667 \pm 0.999	0.71	71.08	2.980 \pm 0.141	0.25	24.95
Diet (5)	38.000 \pm 0.661	0.94	94.22	5.137 \pm 0.337	0.43	43.01
Diet (6)	29.333 \pm 0.366	0.73	72.73	6.757 \pm 0.344	0.57	56.58
Diet (7)	18.333 \pm 0.717	0.45	45.45	12.680 \pm 0.788	1.06	106.17
Control	40.333 \pm 1.001	1.00	100.00	11.943 \pm 0.618	1.00	100.00

Diet (1): Antibiotic (Sodium Sulphate demidin) Diet (2): extract of Thyme Diet (3): extract of Clove Diet (4): extract of Cinnamon Diet (5): extract of black seeds Diet (6): extract of Anise Diet (7): Ripe milk Diet (8): Control

A*= Enzyme activity for diet no. () / Enzyme activity for control * % = percentage relative to control

The data resulted for the biochemical assay summarized in table (2), from Figure 1 the changes in β -glucosidase activity in the homogenate of honeybee, *Apis mellifera* show the low activity in all types of different feeding, Antibiotic (Sodium Sulphate demidin), diet contain extract of Thyme, diet contain extract of Clove, diet contain extract of Cinnamon, diet contain extract of black seeds and diet contain extract of Anise (23.95 %, 31.71%, 34.77%,

24.95%, 43.01% and 56.58%, respectively). On the contrary, the diet containing Ripe milk show high activity (106.17%). The enzyme activity decreases as the glucose chain length increases (Bisaria and Mishra, 1989; Kubicek et al., 1993). That means the increase in activity of β -glucosidases due to degradation of diet polysaccharides to furnish monosaccharide units, which are then able to be metabolically and convert monosaccharide to energy (Murray et al., 1994). The activity of the enzyme depends on several factors including; age of bee, stage of the colony, nectar flow, environment conditions and the beekeeping practices β -glycosidase has been purified the ventricles and honey sac of *Apis mellifera* L. (De Moraes and Bowen, 2000).




Fig. 1. Enzyme activity of β - glucosidase (ug glucose./min/mg) on honey bee with different feeding

From Table 2 and Figure 2, The results showed that, the lactate dehydrogenase activity showed highly enzyme activity in diet contain extract of Clove and Antibiotic ((Sodium Sulphate demidin) 215.70% and 192.56%, respectively. Lactate dehydrogenase is an important glycolytic enzyme present in virtually all animal tissues (Kaplan and Pesce, 1996). It is also involved in carbohydrate metabolism and has been used to indicate exposure to chemical stress (Wu and Lam, 1997; Diamantino *et al.*, 2001). Higher LDH activity in insects is most probably due to consumption as well as utilization of large quantities of food and involved in the production of energy, being particularly important when a considerable amount of additional energy is required immediately. A negative correlation between LDH activity and ambient oxygen levels (Wu and Lam, 1997; Diamantino *et al.*, 2001; Senthil Nathan *et al.*, 2005). The result in other diet used in this study show low activity of LDH in the diet contain extract of Thyme, diet contain extract of Cinnamon, diet

contain extract of black seeds , diet contain extract of Anise and diet containing Ripe milk (79.34%, 71.08%, 94.22%, 72.73% and 45.45% , respectively). From result found inverse relationship between β -glucosidase activity and Lactate dehydrogenase activity in diet contain Ripe milk that may be come back to increase of protein percentage and the diets which content antibiotic and ripe milk recorded higher value percentage to area of sealed worker brood reared, mean number of combs covered with bees and pollen stored so that recommended to use that diet for feeding honey bee.




Fig. 2. Enzyme activity of lactate dehydrogenase (U x 10³ /ml) on honey bee with different feeding.

References

- Arrese, E. L. and Soulages, J. I. (2010). Insect fat body: energy, etabolism, and regulation. Annual Review of Entomology 55:207-225.
- Beenakkeras, M. Th. (1969). Carbohydrate and fat as a fuel for insect flight. A comparative study. Journal of Insect Physiology 15:353-361.
- Bishop, G. H. (1925). Body fluid of the honey bee larva. II. Chemical constituents of the blood, and their osmotic effects. Journal of Biological Chemistry 66:77-88.
- Bounias, M. and Morgan, M. R. J. (1990). Effect of sucrose feeding on the induction of honey bee haemolymph alpha glucosidase (E.C.3.2.1.20). Journal of Apicultural Research 29:181-186.
- Candy, D. J. (1989). Utilization of fuels by the flight muscles. Insect flight. CRC Press, Boca Raton, Florida. pp. 305–319.
- Chipoulet, J. M., Chararas, C. (1985). Survey and electrophoretical separation of the glycosidases of *Rhagium inquisitor* (Coleoptera: Cerambycidae) larvae Comparative Biochemistry and Physiology Part B: Comparative Biochemistry 80:241-246.
- De Moraes, R. L. S., and Bowen, I. D. (2000). Modes of cell death in the hypopharyngeal gland of the honey bee (*Apis mellifera* L). Cell Biology International 24:737-743.

- Deutsche Gesellschaft für Klinische Chemie (1972). Empfehlung der Deutschen Gesellschaft für Klinische Chemie (DGCK). *Journal of Clinical Chemistry and Clinical Biochemistry*. 10:182-193.
- Diamantino, T. C., Amadeu, E., Soares, M. V. M., Guilhermino, L. (2001). Lactate dehydrogenase activity as an effect criterion in toxicity tests with *Daphnia magna* Straus. *Chemosphere* 45:553–560.
- Esen, A. (1993). β -glucosidases: overview. β -Glucosidases: Biochemistry and Molecular Biology. American Chemical Society 533. Washington DC. pp. 1–14.
- Ferreira, C., Parra, J. P. P., Terra, W. R. (1997). The effect of dietary plant glycosides on larval midgut β -glucosidases from *Spodoptera frugiperda* and *Diatraea saccharalis*. *Insect Biochemistry and Molecular Biology* 27:55–59.
- Ferreira, C., Torres, B. B., Terra, W. R. (1998). Substrate specificities of midgut β -glucosidases from insects of different orders. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* 119:219–225.
- Feuerbacher, E., Fewell, J. H., Roberts, S. P., Smith, E. F. and Harrison, J. F. (2003). Effects of load type (pollen or nectar) and load mass on hovering metabolic rate and mechanical power output in the honey bee *Apis mellifera*. *Journal of Experimental Biology*, 206:1855-1865.
- Ghadamyari, M., Hosseini-Navah, V., Sharifi, M. (2010). Partial biochemical characterization of α - and β -glucosidases of lesser mulberry pyralid, *Glyphodes pyloalis* Walker (Lep.: Pyralidae). *Comptes rendus biologies* 333:197-204.
- Harrison, J. F., and Fewell, J. H. (2002). Environmental and genetic influences on flight metabolic rate in the honey bee, *Apis mellifera*. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 133:323-333.
- Hrassing, N. and Crailsheim, K. (2005). Differences in drone and worker physiology in honeybees (*Apis mellifera*). *Apidologie* 36:255-277.
- Huber, R. E. and Mathison, R. D. (1976). Physical, chemical and enzymatic studies on the major sucrose on honey bees (*Apis mellifera*). *Canadian Journal of Biochemistry* 54:153–164.
- Kaper, T., Lebbink, J. H. G., Pouwels, J., Kopp, J., Schulz, G. E., van der Oost, J., and de Vos, W. M. (2000). Comparative structural analysis and substrate specificity engineering of the hyperthermostable β -glucosidase CelB from *Pyrococcus furiosus*. *Biochemistry* 39:4963-4970.
- Kaplan, L. A. and Pesce, A. J. (1996). *Clinical Chemistry: Theory, Analysis, Correlation*, third edition. Mosby-Year Book Inc., St. Louis. pp. 609–610.
- Kubicek, C. P., Messner, R., Gruber, F., Mach, R. L. and Kubicek-Pranz, E. M. (1993). The *Trichoderma* cellulase regulatory puzzle: from the interior life of a secretory fungus. *Enzyme and Microbial Technology* 15:90-99.
- Lindorth, R. L. (1988). Hydrolysis of phenolic glycosides by mid gut β -glucosidases in *Papilio glaucus* subspecies. *Insect Biochemistry* 18:789-792.
- Marana, H. R. C., da Silva, J. S. and de Andrade, J. M. (2000). NK cell activity in the presence of IL-12 is a prognostic assay to neoadjuvant chemotherapy in cervical cancer. *Gynecologic oncology* 78:318-323.
- Mattiacci, L., Dicke, M., and Posthumus, M. A. (1995). β -Glucosidase: an elicitor of herbivore-induced plant odor that attracts host-searching parasitic wasps. *Proceedings of the National Academy of Sciences* 92:2036-2040.
- Murray, R. K. and Granner, D. K., Mayes, P. A. and Rodwell, V. W. (1994). *Harper: Bioquímica*; Atheneu: São Paulo. pp. 628–646.

- Pratviel-Sosa, F., Clermont, S., Percheron, F., and Chararas, C. (1987). Studies on glycosidases and glucanases in *Thaumetopoea pityocampa* larvae—II. Purification and some properties of a broad specificity β -d-glucosidase. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry* 86:173-178.
- Riseh, N. S., Ghadamyari, M. and Motamediniya, B. (2012). Biochemical characterization of α and β -glucosidases and α and β -galactosidases from red palm weevil, *Rhynchophorus ferrugineus* Olivier (Col:Curculionidae). *Plant Protection Science* 2:85–93.
- Rothe, U. and Nachtigall, W. (1989). Flight of the honeybee. IV. Respiratory quotients and metabolic rates during sitting, walking and flying. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry* 158:739–749.
- Sacktor, B. (1970). Regulation of intermediary metabolism, with special reference to the control mechanisms in insect flight muscle. *Advances in Insect Physiology* 7:267-347.
- Santos, C. D. and Terra, W. R. (1985). Physical properties, substrate specificities and a probable mechanism for a β -D-glucosidase (cellobiase) from midgut cells of the cassava hornworm (*Erinnyis ello*). *Biochimica et Biophysica Acta (BBA)-Protein Structure and Molecular Enzymology* 831:179-185.
- Senthil Nathan, S., Chung, P. G. and Murugan, K. (2006). Combined effect of biopesticides on the digestive enzymatic profiles of *Cnaphalocrocis medinalis* (Guenée) (the rice leaffolder) (Insecta: Lepidoptera: Pyralidae). *Ecotoxicology and environmental safety* 64:382-389.
- Senthil Nathan, S., Kalaivani, K., Chung, P. G. and Murugan, K. (2006a). Effect of neem limonoids on lactate dehydrogenase (LDH) of the rice leaffolder, *Cnaphalocrocis medinalis* (Guenee) (Insecta: Lepidoptera: Pyralidae). *Chemosphere* 62:1388–1393.
- Senthil Nathan, S., Kalaivani, K. and Murugan, K. (2006b). Effect of biopesticides on the lactate dehydrogenase (LDH) of the rice leaffolder, *Cnaphalocrocis medinalis* (Guenee) (Insecta: Lepidoptera: Pyralidae). *Ecotoxicology and environmental safety* 65:102–107.
- Sharifi, M., Ghadamyari, M., Moghadam, M. M., and Saïdi, F. (2011). Biochemical characterization of digestive carbohydrases from *Xanthogaleruca luteola* and inhibition of its α -amylase by inhibitors extracted from the common bean. *Archives of Biological Sciences* 63:705-716.
- Tanimura, T., Kitamura, K., Fukuda, T. and Kikuchi, T. (1979). Purification and partial characterization of three forms of alpha-glucosidase from the fruit fly *Drosophila melanogaster*. *Journal of Biochemistry* 1:123–130.
- Terra, W. R. and Ferreira, C. (1983). Further evidence that enzymes involved in the final stages of digestion by *Rhynchosciara* do not enter the endoperitrophic space. *Insect Biochemistry* 2:143–150.
- Terra, W. R. and Ferreira, C. (1994). Insect digestive enzymes: properties, compartmentalization and function. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry* 109:1-62.
- Wheeler, C. H. (1989). Mobilization and transport of fuels to the flight muscles. *Insect flight*. CRC Press, Boca Raton, Florida. pp. 273–303.
- Wu, R. S. S. and Lam, P. K. S. (1997). Glucose-6-phosphate dehydrogenase and lactate dehydrogenase in the green-lipped mussel (*Perna viridis*). Possible biomarker for hypoxia in the marine environment. *Water Research* 31:2797–2801.
- Zidan, E. W. (2009). Studies on varroa mite and its effect on productivity of honeybee colonies. (Doctor of Philosophy's Thesis). Benha University.

(Received 31 January 2014; accepted 28 February 2014)