

---

## Three New Species of Cercosporoid Fungi from Northern Thailand

---

Wongsopa, K. and Cheewangkoon, R.\*

Department of Entomology and Plant Pathology, Faculty of Agriculture, Chiang Mai University, 50200, Thailand.

Wongsopa, K. and Cheewangkoon, R. (2015). Three new species of Cercosporoid fungi from Northern Thailand. *International Journal of Agricultural Technology* 11(1):127-141.

**Abstract** A collection of Cercosporoid fungi occurring in leaf spots from Northern Thailand during 2013–2014 is described and illustrated. These fungi include three new species, *Cercospora posoneae*, *Pseudocercospora combretina* and *Pseudocercospora glycinicola*. Eleven species are new records of host plants for Thailand. Six species of *Cercospora* morphologically similar to *Cercospora apii* sensu lato were also found in this study.

**Keywords:** Cercosporoid, *Cercospora*, *Pseudocercospora*

### Introduction

Cercosporoid fungi are well known as one of the largest hyphomycete genera, and numerous species of this fungus have been linked to leaf spots symptoms. Many hyphomycetes genera are linked to fungus *Mycosphaerella* and many host plant species are damaged by these fungi (Crous, 2009).

Among the Cercosporoid fungi, *Cercospora* and *Pseudocercospora* genera are considered to be important fungi because of numerous plant disease reports from Northern Thailand (Gavin *et al.*, 2006; Nakachima *et al.*, 2007; Meeboon *et al.*, 2007; To-anun *et al.*, 2011). They cause lesion symptoms, including necrotic leaf spots and leaf blight on many host plants including as agronomic crops, ornamental plants and forest trees (Crous *et al.*, 2004, 2006, 2007; Arzanlou *et al.*, 2008).

Ecologically these fungi are also known to be saprobes, parasites of other fungi (Jackson *et al.*, 2004) as well important plant pathogens, and are mostly found in tropical and sub-tropical areas. Although, most *Cercospora*

---

\* **Corresponding author:** Cheewangkoon, R.; **Email:** [ratcha.222@gmail.com](mailto:ratcha.222@gmail.com)

species are host specific (Arzanlou *et al.*, 2007), but some species still produce lesions on various different plants (Groenewold *et al.*, 2006) through co-infection with an unrelated fungus in the same lesion.

The present study on Cercosporoid fungi is similar to those conducted in many other countries which have sought to discover new *Cercospora* species. Moreover, there have been numerous reports of new species, new hosts and new records of Cercosporoid fungi in South–East Asia (Burgess *et al.*, 2007). In Thailand and neighboring countries there have been many reports of new Cercosporoid fungi, such as *Cercospora*, *Passalora*, *Pseudocercospora* and other fungi. (Nakachima *et al.*, 2007; Meeboon *et al.*, 2007; To-anun *et al.*, 2011; Phengsintham *et al.*, 2013). These reports were mainly based on the morphological taxonomic concepts introduced by Chupp (1954) and further developed and modified by Crous & Braun (2003). The taxonomic identification key of Cercosporoid fungi used in the current study was based on these reports.

## Materials and methods

The specimens were collected in Chiang Mai, Chiang Rai and Lampang provinces in Northern Thailand. Leaf lesions were examined under a stereo microscope for the presence the Cercosporoid fungi. The slides for microscopic examination were prepared by hand sections from plant materials and mounted in lactic acid.

Detailed observations of morphological characteristics were carried out by means of an Olympus CX 31 light microscope (100×). Thirty conidia, conidiophores and stromata were measured for each specimen. Dried specimens were deposited at Plant Pathology Herbarium, Department of Agriculture, Chiang Mai University.

## Results

*Cercospora pisoneae* Wongsopa and Cheewangkoon, sp. nov.

*Etymology*: pisoneae, derived from the genus name of the host plant. Maculae amphigenae, circulares vel subcirculares, centro griseo-brunnea, margine indefinitis, 5 mm diam. Caespituli amphigeni. Stromata nulla vel parva, evolutis, substomatalia, 10–30 µm diam., atro-brunnea. Conidiophora 3–5 laxe fasciculate, cylindrical, erecta vel decumbentia, recta vel leviter curvata, 1–3-geniculata, non ramosa, (45–) 275–370 (–400) × 4–7


$\mu\text{m}$ , 8–10 (–13) –septata, medio brunnea, laevia. Cellulae conidiogena terminalis, sympodiales, Loci conidiogeni conspicui, incrassati, fuscati. Conidia solitaria, fusiformia vel obclavata, recta vel leviter curvata, (200–) 215–340 (–380)  $\times$  (3–) 4 (–5)  $\mu\text{m}$ , (16–) 17–28 (–32) –septata, hyaline, non ramosa, laevia, apice subacuto vel obtusa et basi obconice truncata, hila incrassata et fuscata.

*Leaf spots* circular to subcircular, centre grayish to pale brown and dark brown indefinite margins, 5 mm in diameter. *Caespituli* amphigenous. *Stromata* lacking to small–developed, subepidermal, arising from stomata cells, 10–30  $\mu\text{m}$  in diameter, dark brown. *Conidiophores* loosely fasciculate, arising from stromata (3–5 per fascicle), cylindrical, erect to decumbent, straight or slightly curved, sometimes 1–3–geniculate at base, unbranched, (45–) 275–370 (–400)  $\times$  4–7  $\mu\text{m}$ , 8–10 (–13) –septate, medium brown, smooth. *Conidiogenous cells* terminal, conidiogenous loci conspicuous, thickened and darkened. *Conidia* solitary, filiform to narrowly obclavate, straight to slightly curved, (200–) 215–340 (–380)  $\times$  (3–) 4 (–5)  $\mu\text{m}$ , (16–) 17–28 (–32) –septate, hyaline, unbranched, smooth, apex obtuse and base obconically truncated, *Hila* thicken and darkened. (Fig. 1–2)


*Habitat*: On leaves of *Pisonia grandis* R. Br. (Nyctaginaceae)

*Material examined*: Thailand, Chiang Mai Province, Mae Rim district, Nang Hoi, 26 July 2013, Khelang Wongsopa and Ratchadawan Cheewangkoon (Plant Pathology Herbarium HY32).

*Notes*: Numerous reports can be found describing *Cercospora* spp. on Nyctaginaceae, but this is the first report of *Cercospora pisoneae* on Genus *Pisonea* (Nyctaginaceae) from Thailand.



**Figure 1.** *Cercospora pisoneae* A. Leaf spots on *Pisonia grandis* R. Br. (Scale bars = 1 cm) B. Caespituli, C–D. Stromata and conidiophores, E. Conidia (C–E: Scale bars = 10  $\mu\text{m}$ ).



**Figure 2.** *Cercospora pisoneae* on *Pisonia grandis* R. Br. (From holotype). A. Conidia B. Stromata and conidiophores. Scale bars = 50  $\mu$ m.

***Pseudocercospora combretina*** Wongsopa and Cheewangkoon, sp. nov.

*Etymology:* combretina, derived from the genus name of the host plant. Maculae amphigenae subcirculares vel irregulares, centro griseo–albidis vel pallide brunnea, medio brunneo vel atro–brunnea margine indefinitis, 5–15 mm diam., Caespituli epiphyllis. Stromata bene–evolūtis, substomatalia vel


intraepidermalia, usque 50  $\mu\text{m}$  dium., subglobosis, atro-brunnea vel cinereo. Conidiophora 15–25 laxe vel dense fasciculate, aciculis vel subcylindrica, erecta vel decumbentia, recta vel leniviter curvata, geniculate et apice, non ramosa, (6–) 10–20  $\times$  2.5–4  $\mu\text{m}$ , 0–1 (–2) -septata, pallide vel medio brunneo. Laevia, rotundata vel subconico et apice. Cellulae conidiogenae terminalis. Loci conidiogeni inconspicui. Conidia solitaria, obclavata vel subcylindrica, recta vel leniviter curvata, sinuosa, (12–) 20–82 (–88)  $\times$  3–4  $\mu\text{m}$ , 1–5-septata, conspicui, brunneo ad apicem versus pallidiora, non ramosa, laevia, apice obtuse, et basi obconice truncata.

*Leaf spots* subcircular to irregular, centre grayish, white to pale brown and medium brown to dark brown indefinite margins, 5–15 mm in diameter. *Caespituli* mainly epiphyllous. *Stromata* well-developed, subepidermal to intraepidermal, up to 50  $\mu\text{m}$  in diameter, subglobular, dark brown to blackish. *Conidiophores* loosely to moderate fasciculate, arising from stromata (15–25 per fascicle), acicular to subcylindrical, erect to decumbent, straight or slightly curved, sometimes geniculate at apex, unbranched, (6–) 10–20  $\times$  2.5–4  $\mu\text{m}$ , 0–1 (–2) -septate, pale to medium brown, smooth, small rounded to subconic apex. *Conidiogenous cells* terminal, conidiogenous loci inconspicuous, unthickened and not darkened. *Conidia* solitary, narrowly obclavate to subcylindrical, straight to slightly curved, sinuous, (12–) 20–82 (–88)  $\times$  3–4  $\mu\text{m}$ , 1–5-septate conspicuous, brown, and paler toward the apex, unbranched, smooth, apex obtuse and base obconically truncated. (Fig. 3–4)


*Habitat*: On leaves of *Combretum procursum* Craib (Combretaceae)

*Material examined*: Thailand, Lampang Province, Chaeson National Park, 8 January 2014, Khelang Wongsopa and Ratchadawan Cheewangkoon (Plant Herbarium HY41).

*Notes*: This is a new record for this species from Thailand. Previously two species of *Pseudocercospora* spp. were reported on *Combretum* spp. The current specimen is distinguished by having *stromata* dark brown to blackish, *Conidiophores* are shorter (6–) 10–20  $\times$  2.5–4  $\mu\text{m}$  and *Conidia* are brown to dark brown.



**Figure 3.** *Pseudocercospora combretina* A. Leaf spots on *Combretum procursum* Craib (Scale bars = 1 cm), B. Caespituli, C. Stromata and conidiophores, D. Conidia (C–D: Scale bars = 10  $\mu\text{m}$ ).



**Figure 4.** *Pseudocercospora combretina* on *Combretum procursum* Craib (From holotype). A. Conidia, B. Stromata and conidiophores. (Scale bars: = 50  $\mu\text{m}$ .)

***Pseudocercospora glycinicola* Wongsopa & Cheewangkoon, sp. nov.**

*Etymology*: glycinicola, derived from the genus name of the host plant.


Maculae amphigenae, subcirculares vel irregulares, griseo vel brunneo, 1–5 mm dium., Caespituli amphigenae. Stromata nulla vel parva–evolutis, substomatalia, 20–25 (–50)  $\mu\text{m}$  diam., brunneo vel atro-brunnea, Conidiophora 15–21 fasciculata, subcylindrica, erecta vel decumbentia, recta, leviter curvata vel sinuosa, ramosa, (30–) 40–70 (–75)  $\times$  5–6 (–7)  $\mu\text{m}$ , 4–6-septata, laevia, pallide brunnea vel brunneo, ad apicum versus pallidiora, apicem subacuto vel rotundato ad subtruncata. Cellulae conidiogenae terminalis, sympodiale, Loci conidiogeni inconspicui. Conidia solitaria, subcylindrica vel cylindrical, obclavata, recta vel leviter curvata, (40–) 65–95 (–100)  $\mu\text{m}$ , (3–) 4–7-septata, guttulata, subhyalina vel pallide brunnea, laevia, apice obtuse vel rotundato, et basi obconice truncate.

*Leaf spots* subcircular to irregular, grayish to brown, 1–5 mm in diameter, *Caespituli* amphigenous. *Stromata* lacking to small–developed, substomatal, 20–25 (–50)  $\mu\text{m}$  in diameter, brown to dark brown. *Conidiophores* fasciculate (15–21 per fascicle), erumpent through the cuticle, subcylindrical and narrower towards at apex, erect to decumbent, straight, slightly curved to sinuous, branched, (30–) 40–70 (–75)  $\times$  5–6 (–7)  $\mu\text{m}$ , 4–6-septate, smooth, brown to pale brown, and paler toward at apex, subacute to round and subtruncate at apex. *Conidiogenous cells* sympodial, terminal, conidiogenous loci inconspicuous, unthickened, not darkened. *Conidia* solitary, subcylindrical to cylindrical, sometimes narrowly obclavate, mostly straight to slightly curved, (40–) 65–95 (–100)  $\mu\text{m}$ , (3–) 4–7-septate, guttulate, subhyaline to pale brown, smooth, apex obtuse, base obconically truncated. (Fig. 5–6)


*Habitat*: On leaves of *Glycine max* (L.) Merr. (Fabaceae)

*Material examined*: Thailand, Chiang Mai Province, Mae Wang district, 26 July 2013, Khelang Wongsopa (Plant Pathology Herbarium HY25).

*Notes*: Only *Pseudocercospora glycines* species found on Genus *Glycine* spp. by Hsieh and Goh (1990), *P. glycines* was previously reported only from Australia and Taiwan. *P. glycinicola* is distinguished by having *conidiophores* branched and longer than *P. glycines* (30–) 40–70 (–75)  $\times$  5–6 (–7)  $\mu\text{m}$  Vs. 10–25  $\times$  1.5–3  $\mu\text{m}$ .



**Figure 5.** *Pseudocercospora glycinicola* A. Leaf spots on *Glycine max* (L.) Merr (Scale bars = 1 cm), B. Caespituli, C. Stromata and conidiophores, D. Conidia (C–D: Scale bars = 10  $\mu\text{m}$ ).



**Figure 6.** *Pseudocercospora glycinicola* on *Glycine max* (L.) Merr. (From holotype). A. Conidia B. Stromata and conidiophores. Scale bars: = 50  $\mu\text{m}$ .

***Cercospora apii*** Fresen. *sensu lato*, emend. Crous and Braun, *Mycosphaerella* and its anamorphs: 1. Names published in *Cercospora* and *Passalora*: 35. 2003.

Crous and Braun (2003) organized the several phylogenetic species and their morphologically similar species of the *Cercospora apii* complex from many host species. However, their scheme lacks the taxonomic identification of the genus *Cercospora* from Northern Thailand.

***Cercospora apii*** (Crous and Braun, 2003)

*Caespituli* amphigenous. *Stromata* substomatal to intraepidermal, 25–40 µm in diam., dark brown. *Conidiophores* cylindrical, 125–240 (–250) × 5–6 µm, (2–) 3–8 (–9) -septate, brown. *Conidia* filiform to narrowly obclavate, (35–) 70–300 (–380) × (3–) 4–5 µm, (2–) 6–22 (–29) -septate, hyaline to subhyaline.

*Habitat*: On leaves of *Brassica rapa* L. (Cruciferae)

*Material examined*: Thailand, Chiang Mai Province, Multiple Cropping Center, Faculty of Agriculture Chiang Mai University, 15 August 2013, Khelang Wongsopa and Ratchadawan Cheewangkoon (Plant Pathology Herbarium HY33).

*Notes*: This is the first report of a new host from Thailand.

***Cercospora ipomoeae*** (Hsieh and Goh, 1990)

(=*Cercospora apii* Crous and Braun, 2003)

*Caespituli* amphigenous, *Stromata* substomatal, 20–60 µm in diam., dark brown to blackish. *Conidiophores* cylindrical, (45–) 50–120 × 5–6 µm, 2–3-septate, unbranch, pale brown to medium brown. *Conidia* narrowly obclavate, (75–) 100–210 (–310) × 4–5 µm, (7–) 10–15 (–17) -septate, guttulate, hyaline to subhyaline.

*Habitat*: On leaves of *Merremia vitifolia* (Burm. f.) Hallier f. (Convolvulaceae)

*Material examined*: Thailand, Chiang Mai Province, Mueang District, Mae Hia, 10 June 2013, Khelang Wongsopa and Ratchadawan Cheewangkoon (Plant Pathology Herbarium HY20).

*Notes*: This is the first report of this species on a new host from Thailand.

***Cercospora apii*** (Crous and Braun, 2003)

*Caespituli* amphigenous, *Stromata* substomatal, 25–80 µm in diam., dark brown. *Conidiophores* cylindrical, (80–) 85–180 (–190) × (4–) 5–6 (–7)

$\mu\text{m}$ , (2–) 3–4 (–5)-septate, unbranch, pale brown. *Conidia* narrowly obclavate–subcylindrical, (70–) 90–275 (–300)  $\times$  4–5  $\mu\text{m}$ , (8–) 13–21 (–24)-septate, hyaline.

*Habitat*: On leaves of *Abelmoschus esculentus* (L.) Moench (Malvaceae)

*Material examined*: Thailand, Chiang Mai Province, Mae Wang District, 28 June 2013, Khelang Wongsopa and Ratchadawan Cheewangkoon (Plant Pathology Herbarium HY22).

*Notes*: This is the first report on this host from Thailand.

***Cercospora apii*** (Crous and Braun, 2003)

*Caespituli* amphigenous, *Stromata* substomatal, 50–65  $\mu\text{m}$  in diam., dark brown. *Conidiophores* cylindrical, (130–) 140–190 (–210)  $\times$  5–6  $\mu\text{m}$ , 4–6 (–8) -septate, unbranch, pale brown. *Conidia* filiform–narrowly obclavate, 70–280  $\times$  4  $\mu\text{m}$ , 10–28-septate, gutulate, hyaline.

*Habitat*: On leaves of *Thunbergia laurifolia* Lindl. (Acanthaceae)

*Material examined*: Thailand, Chiang Mai Province, Muang District, Mae Hia, 14 February 2013, Khelang Wongsopa and Ratchadawan Cheewangkoon (Plant Pathology Herbarium HY15).

*Notes*: This is the first report of a new host from Thailand.

***Cercospora paederiicola*** (Guo and Liu, 1991)

(=*Cercospora apii* Crous and Braun, 2003)

*Caespituli* amphigenous, *Stromata* substomatal, 100–120  $\mu\text{m}$  in diam., dark brown to blackish. *Conidiophores* cylindrical, (110–) 115–130 (–150)  $\times$  4–6  $\mu\text{m}$ , 5–6-septate, unbranch, pale brown. *Conidia* narrowly obclavate, (75–) 100–210 (–310)  $\times$  4–5  $\mu\text{m}$ , (11–) 13–25 (38) -septate, gutulate, hyaline.

*Habitat*: On leaves of *Paederia pallida* Craib (Rubiaceae)

*Material examined*: Thailand, Chiang Mai Province, Muang District, Mae Hia, 10 June 2013, Khelang Wongsopa and Ratchadawan Cheewangkoon (Plant Pathology Herbarium HY18).

*Notes*: This is the first report of a new host from Thailand.

***Pseudocercospora liquidam Scalebarsicola*** (J.M. Yen) Braun, (2000)

*Caespituli* amphigenous, *stromata* substomatal, up to 65 in diam., dark brown. *Conidiophores* 9–10 (–20)  $\times$  3–4  $\mu\text{m}$ , 0–1-septate, unbranched, pale brown. *Conidia* acicular–narrowly obclavate, subcylindrical (30–) 35–52 (–65)  $\times$  (2.5–) 3  $\mu\text{m}$ , (2–) 3–5 (–7) -septate, gutalate, greyish to brown.

*Habitat:* On leaves of *Liquidambar formosana* Hance. (Hamamelidaceae)

*Material examined:* Thailand, Chiang Mai Province, Chom Thong District, Ban Mae Tao, 13 November 2012, KhelangWongsopa and RatchadawanCheewangkoon (Plant Pathology Herbarium HY1).

*Notes:* This is the first report of this species from Thailand.

***Pseudocercospora fici*** Heald & F.A. Wolf (Liu and Guo, 1991)

(=*Cercospora fici* Heald & Wolf, 1911)

*Caespituli* amphigenous. *Stromata* substomatal, 25–35  $\mu\text{m}$  in diam, brown to dark brown. *Conidiophores* (9–) 10–20 (–25)  $\times$  (2.5–) 3–3.5 (–4)  $\mu\text{m}$ , 0–1-septate, unbranched, subhyaline to pale brown. *Conidia* subcylindrical, (60–) 65–80  $\times$  (2.5–) 3 (–4)  $\mu\text{m}$ , (4–) 5–8 (–10) -septate, gutulate, subhyaline to pale brown.

*Habitat:* On leaves of *Ficus pumila* L. (Moraceae)

*Material examined:* Thailand, Chiang Rai Province, Mae Fah Luang District, Doi Tung, 3 February 2013, Khelang Wongsopa and Ratchadawan Cheewangkoon (Plant Pathology Herbarium HY9).

*Notes:* This is the first report of this species on a new host from Thailand.

***Pseudocercospora mori*** (Hara) Deighton, (1976)

*Caespituli* hypophyllous, *Stromata* substomatal, 35–55  $\mu\text{m}$  in diam., dark brown. *Conidiophores* cylindrical, (5–) 6–9 (–10)  $\times$  2–5  $\mu\text{m}$ , aseptate, unbranched, pale brown. *Conidia* narrowly obclavate–subcylindrical, (20–) 25–55 (–65)  $\times$  2–3  $\mu\text{m}$ , (1–) 2–6 (–7) -septate, gutulate, *subhyaline to greyish*.

*Habitat:* On leaves of *Morus alba* L. (Moraceae)

*Material examined:* Thailand, Chiang Mai Province, Muang District, Faculty of Agriculture, Chiang Mai University, 15 August 2013, Khelang Wongsopa and Ratchadawan Cheewangkoon (Plant Pathology Herbarium HY34).

*Notes:* This is the first report of this species on a new host from Thailand.

***Pseudocercospora rubi*** (Sacc.) Deighton, (1976)

*Caespituli* epiphyllous, *Stromata* substomatal–intraepidermatal, 40–80  $\mu\text{m}$  in diam., dark brown. *Conidiophores* cylindrical, geniculate–sinuous,

(10–) 15–20 (–25) × 3–4 μm, 0–1-septate, unbranched, pale brown to hyaline. *Conidia* mostly curved, subcylindrical–cylindrical, (40–) 50–80 (–105) × 3–4 μm, (1–) 3–8 (–9) -septate, gutulate, grayish to subhyaline.

*Habitat:* On leaves of *Rubus blepharoneurus* Card. (Rosaceae)

*Material examined:* Thailand, Chiang Rai Province, Thoeng District, Phu Chi Fa, 10 December 2013, Khelang Wongsopa and Ratchadawan Cheewangkoon (Plant Pathology Herbarium HY38).

*Notes:* This is the first report of this species on a new host from Thailand.

***Pseudocercospora egenula* (Syd.) Braun and Crous, (2003)**

*Caespituli* amphigenous, *Stromata* substomatal, 25–45 μm in diam., brown to dark brown. *Conidiophores* subcylindrical, (17–) 20–35 (–45) × (3.5–) 4–5 (–6) μm, (0–) 1–2 (–4) -septate, unbranched, pale brown to brown. *Conidia* narrowly obclavate–subcylindrical, (16–) 35–75 (–95) × 3–4 (–5) μm, 1–7 (–9) -septate, greyish to subhyaline.

*Habitat:* On Leaves of *Brugmansia × candida* Pers. (Solanaceae)

*Material examined:* Thailand, Chiang Mai Province, Chom Thong District, Doi Intanon, 27 January 2013, Khelang Wongsopa and Ratchadawan Cheewangkoon (Plant Pathology Herbarium HY4).

*Notes:* This is the first report of this species on a new host from Thailand.

***Pseudocercospora daturina* (J. M. Yen) Deighton, (1976)**

*Caespituli* amphigenous, *Stromata* substomatal, 35–65 μm in diam., brown to dark brown. *Conidiophores* subcylindrical, (10–) 15–40 (–45) × 4–5 (–6) μm, 0–2 (–3) -septate, branched to unbranched, greyish to pale brown. *Conidia* narrowly obclavate to subcylindrical, (22–) 42–87 (–110) × (3–) 4–5 μm, (1–) 4–8 (–11) -septate, gutulate, grayish to pale brown.

*Habitat:* On Leaves of *Brugmansia × candida* Pers. (Solanaceae)

*Material examined:* Thailand, Chiang Rai Province, Mae Fah Luang District, Doi Tung, 3 February 2013, Khelang Wongsopa and Ratchadawan Cheewangkoon (Plant Pathology Herbarium HY10).

*Notes:* This is the first report of this species on a new host from Thailand.

## Discussion

The Cercosporoid fungi are well known to be associated with *Mycosphaerella* teleomorphic fungi, and they have been linked to leaf spots symptoms on many hosts. Numerous reports can indicate that they include some of most important living tissue pathogens in tropical regions (Crous, 2009).

A specific focus of this study was to survey for Cercosporoid fungi in forests and plantations in Northern Thailand, because a number of previous reports on these fungi identified new species, new hosts and new records (Meeboon *et al.*, 2007; Nakachima *et al.*, 2007; To-anun *et al.*, 2011; Phengsintham *et al.*, 2011).

Our research obtained 14 isolates of *Cercospora* spp. and *Pseudocercospora* spp. from lesions on host plants, including new species from Thailand: *Cercospora pisoneae* on *Pisonia grandis* (Nyctagina), *Pseudocercospora combretina* on *Combretum procursum* Craib (Combretaceae) and *Pseudocercospora glycinicola* on *Glycine max* (L.) Merr (Fabaceae). These isolates differed taxonomically from all known Cercosporoid fungi and can be considered new species.

In addition, the lists of *Cercospora apii sensu lato* on many host plants in this report was consistent reporting in “*Mycosphaerella* and its anamorph: 1. Names published in *Cercospora* and *Passalora*” reported by Crous and Braun (2003).

## Acknowledgements

This work was financially supported by Thailand Research Fund (DBG5380011 and MRG5580163). The authors thank Dr. J. F. Maxwell for identification of the host plants.

## References

- Arzanlou, M., Abeln, E. C. A., Kama, G. H. J., Waalwijk, C., Carlier, J., de Vries, I., guzman, M. and Crous, P. W. (2007). Molecular diagnostics for the Sigatoka disease complex of banana. *Phytopathology* 97:1112-1118.
- Arzanlou, M., Groenewald, J. Z., Fullerton, R. A., Abeln, E. C. A., Carlier, J., Zapater, M. F., Buddenhagen, I. W., Viljoen, A. and Crous, P. W. (2008). Mutiple gene genealogies and phenotypic characters differentiate several novel species of *Mycosphaerella* and related anamorphs on banana. *Persoonia* 20:19-37.

- Braun, U. (2000). Miscellaneous notes on some micromycetes. *Schlechtendalia* 5:31-56.
- Burgess, T. I., Scale barsber, P. A., Sufaati, S., Xu, D., Hardy, G. E. S. and Dell, B. (2007). *Mycosphaerella* spp. on *Eucalyptus* in Asia; new species; new hosts and new records. *Fungal Diversity* 24:135-157.
- Chupp, C. (1954). A monograph of the fungus genus *Cercospora*. New York: Chupp, Charles.
- Crous, P. W. (2009). Taxonomy and phylogeny of the genus *Mycosphaerella* and its anamorphs. *Fungal Diversity* 38:1-24.
- Crous, P. W. and Braun, U. (2003). *Mycosphaerella* and its anamorphs: 1. names published in *Cercospora* and *Passalora*. Utrecht: Centraalbureau voor Schimmelcultures.
- Crous, P. W., Wingfield, M. J., Mandilla, J. P., Alfenas, A. C. and Groenewald, J. Z. (2006). Phylogenetic reassessment of *Mycosphaerella* spp. and their anamorphs occurring on *Eucalyptus*. II. *Studies in Mycology* 55:99-131.
- Crous, P. W., Groenewald, J. Z., Pongpanich, K., Himaman, W., Arzanlou, Z. and Wingfield, M. J. (2004). Cryptic speciation and host specificity among *Mycosphaerella* spp. occurring on Australian *Acacia* species grown as exotics in the tropics. *Studies in Mycology* 50:457-469.
- Crous, P. W., Summerell, B. A., Carnegie A. J., Mohammed, C., Himaman, W. and Groenewald, J. Z. (2007). *Foliicolous Mycosphaerella* spp. and their anamorphs on *Corymbia* and *Eucalyptus*. *Fungal Diversity* 26:143-185.
- Deighton, F. C. (1976). Studies on *Cercospora* and allied genera-VI. *Pseudocercospora* Speg., *Pantospora* Cif. and *Cercoseptoria* petr. *Mycological Papers* 140:1-168.
- Groenewald, M., Groenewald, J. Z., Braun, U. and Crous, P. W. (2006). Host range of *Cercospora* *apii* and *C. beticola* and description of *C. apiicola*, a novel species from celery. *Mycologia* 98:275-285.
- Guo, Y. L. and Liu, X. J. (1991). Studies on the genus *Pseudocercospora* in China V. *Mycosystema* 4:99-148.
- Hsieh, W. H. and Goh, T. K. (1990). *Cercospora* and similar fungi from Taiwan. Taipei: Maw Chang Book Company.
- Hunter, G. C., Crous, P. W., Wingfield, B. D., Pongpanich, K. and Wingfield, M. J. (2006). *Pseudocercospora flavomarginata* sp. nov., from *Eucalyptus* leaves in Thailand. *Fungal Diversity* 22:71-90.
- Jackson, S. L., Maxwell, A., Neumeister-Kemp, H. G., Dell, B. and Hardy G. E. S. J.

- (2004). Infection, hyperparasitism and conidiogenesis of *Mycosphaerella lateralison Eucalyptus globulus* in Western Australia. *Australasian Plant Pathology* 33:49-53.
- Meeboon, J., Hidayat, I. and To-anun, C. (2007). An annotated list of Cercosporoid fungi in Northern Thailand. *Journal of Agricultural of Agricultural Technology* 3:51-63.
- Nakachima, C., Motohashi, K., Meeboon, J. and To-anun, C. (2007). Studies on *Cercospora* and allied genera in Northern Thailand. *Fungal Diversity* 26:257-270.
- Phengsintham, P., Chukeatirote, E., McKenzie, E. H. C., Hyde, K. D. and Braun, U. (2013). Monograph of Cercosporoid fungi from Laos. *Current Research in Environmental and Applied Mycology* 3:34-158.
- To-anun, C., Hidayat, I. and Meeboon, J. (2011) Genus *Cercospora* in Thailand: Taxonomy and Phylogeny (with a dichotomous key to species). *Plant Pathology and Quarantine* 1:11-87.

(Received: 9 November 2014, accepted: 24 December 2014)