

---

## Mycoflora and Mycotoxin Contaminated Some Juices

---

Embaby, E. S. M.<sup>1\*</sup>, Awni, N. M.<sup>2</sup>, Abdel-Galil, M. M.<sup>3</sup> and El-Gendy, H. I.<sup>4</sup>

<sup>1</sup>Plant Pathology Dept., National Research Centre, Cairo, Egypt; <sup>2</sup>Department of Microbiology, Faculty of Science, Zagazig University; <sup>3</sup>Food Toxicology and Contaminants Dept. National Research Centre, Cairo, Egypt; <sup>4</sup>Zefta Hospital, Ministry of Health, Egypt.

Embaby, E. S. M., Awni, N. M., Abdel-Galil, M. M. and El-Gendy, H. I. (2015). Mycoflora and mycotoxin contaminated some juices. International Journal of Agricultural Technology 11(3):693-712.

**Abstract** Fruit juices are popular drinks as they contain antioxidants, vitamins, and minerals that are essential for human being and play important role in the prevention of heart diseases, cancer, and diabetes. Consumption of fresh juices increased dramatically due to their freshness, high vitamin content, and low caloric consumption. Fruit and vegetable markets are known to contain several species of fungi. A total of 60L juice samples were examined for their microbiological association. Ten fungal species including seven fungal genera were isolated and identified from the three types of fresh juices survived i. e. apple, grape-fruit and navel orange. These are *Alternaria citri*, *Aspergillus flavus*, *A. parasiticus*, *Botryodiplodia theobromae*, *Colletotrichum gloeosporioides*, *Fusarium* sp., *Penicillium expansum*, *P. digitatum*, *P. italicum* and *Rhizopus stolonifer*. Tested of mycotoxin production indicated that, *A. flavus* (No. 20) was found to be produced only 0.10 ng/ml of aflatoxin AFB<sub>1</sub>, *A. parasiticus* (No. 4) produced 0.52 ng/ml of aflatoxin AFB<sub>1</sub>, 0.10 ng/ml of aflatoxin AFB<sub>2</sub>, 0.32 ng/ml of aflatoxin AFG<sub>1</sub> and 0.10 ng/ml of aflatoxin AFG<sub>2</sub>. Decontamination of fungi resulted that, all hot water treatments i. e. 45, 50 and 55<sup>0</sup>c were found to reduce the number of spores' viability of the tested fungi (*A. flavus*, *A. parasiticus* and *P. expansum*) compared with un-treated. The numbers of spore viability were decreased with increasing the temperature degree used. Both pasteurization (at 80 °C) and sterilization (at 121 °C) treatments have a great effect on spore viability of the three tested fungi compared with un-treated juices which gave completely inhibition of spore viability of the three tested fungi with hundred percent of reduction.

**Keywords:** Fresh juice, Apple, Grapefruit, Navel orange, Fungi, Mycotoxin

### Introduction

Fruit juices are popular drinks as they contain antioxidants, vitamins, and minerals that are essential for human being and play important role in the prevention of heart diseases, cancer, and diabetes. They contain essential nutrients which support the growth of acid tolerant bacteria, yeasts, and moulds. Consumption of fresh juices increased dramatically due to their freshness, high vitamin content, and low caloric consumption. Extracted juices from fruits

---

\* Corresponding author: Embaby, E. S. M.; Email:-

contain most substances which are found in the original ripe and sound fruit from which the juice is made. The high potassium and low sodium characteristic of most juices help in maintaining a healthy blood pressure. Vitamin C is naturally present in juices which are essential for the body to form collagen, cartilage, muscle, and blood vessels. It also helps in the absorption of iron (Rathnayaka, 2013 and Kamal, *et al.*, 2014).

Apple (*Malus domestica*) is a highly nutritious fruit containing essential food elements such as carbohydrates, protein, fat and water. Apart from its energy value, apple is a good source of soluble and insoluble fiber (Herforth, 2000). Apples are a widely consumed, rich source of phyto-chemicals; epidemiological studies have linked the consumption of apples with reduced risk of some cancers, cardiovascular disease, asthma, and diabetes due to strong antioxidant activity (Marchand *et al.*, 2000).

The most important of the citrus fruits (*Citrus* sp.) (Family Rutaceae or orange family) are commonly eaten fresh, juiced, and in processed products. Citrus fruits are acidic fruits and they are well known for their health promoting compounds. Citrus is a good source of vitamin C (L-ascorbic acid). Citrus fruits also contain an impressive list of other essential nutrients, including glycaemic and non-glycaemic carbohydrates (sugars and fiber), potassium, folate, calcium, thiamin, niacin, vitamin B6, phosphorus, magnesium, copper, riboflavin, pantothenic acid and a variety of phytochemicals. In addition, citrus contains no fat or sodium and, being a plant food, no cholesterol. The rind, which contains numerous oil glands, and the fragrant blossoms of some species are also a source of essential oils used for perfumes and as a flavoring agent in foods. The most important use of citrus for processing is the production of frozen concentrated orange juice (Milind, 2008).

Fruits contain high levels of sugars and nutrients and their low pH values make them vulnerable to fungal decay. It has been estimated that about 20-25% of the harvested fruits are decayed by pathogens during post-harvest handling even in developed countries. Plant pathogens may infect fruits either prior to harvest under field conditions or after harvest during transit and storage. Many pathogens may remain dormant for varying periods until favorable conditions become available for their development, leading to visible symptoms (Droby, 2006; Zhu, 2006; Singh and Sharma 2007). It should be noted that for a total of 100,000 fungi, less than 10% are pathogenic for plants and around 100 species are responsible for the majority of postharvest damage (Singh and Sharma, 2007).

The common postharvest and storage fungi of fruits are *Alternaria* spp., *Aspergillus* spp., *Fusarium* spp., and *Penicillium* spp. (Bhale, 2011). The major mycotoxin-producing fungi are not aggressive pathogens in plants; however,

mycotoxins are produced by several genera in plants during the growing season when portals of entry are provided and environmental conditions are appropriate and be continued or initiated in postharvest and stored products. The majority of these toxins are produced by fungi of the genera, *Aspergillus*, *Penicillium* and *Fusarium* (Barkai-Golan and Paster, 2008 and Ammar and El-Naggar, 2014).

Fruit juices contain a micro-flora which is normally present on the surface of fruits during harvest and postharvest processing which include transport, storage, and processing (Tournas, *et al.*, 2006 and Kamal, *et al.*, 2014). The surface of apple fruits harbors microorganisms depending upon the mechanical handling of the fruits. Microbes can adhere to surface, invade/penetrate fruits surface and multiply within the tissue. Contamination could be from human handling, transport vehicles, insects, dust, and rinse water, harvesting equipment, soil, faeces, irrigation water, water used to apply fungicides and insecticides, manure, wild and domestic animals (Buck *et al.*, 2003). Besides nutritional merits, apple may get contaminated during growth, harvest, transportation and further processing & handling with microbes from soil, air, water or animal wastes. Handling in stores and retail markets could also add more microorganisms to the surface of the fresh produce. Among microbes, *Penicillium expansum*, a fungus is a major causative agent of post harvest decay in apple, since it produces patulin, a mycotoxin known to cause harmful effects in humans (Chen *et al.*, 2004). The mycotoxin patulin is a secondary metabolite produced by different species of fungi in the genera *Penicillium*, *Aspergillus*, *Gymnoascus*, *Paecilomyces* and *Byssosclamyces*. *P. expansum* is the most important patulin producer (Pitt and Hocking, 1997). Although patulin can occur in several fruits, grains and other foods infected by fungi, the main concern are with apples, apple cider and apple juice due to their higher consumption. As mentioned before, patulin can be found in human food and beverages, animal feed, biological and environmental samples (Begum, 2012).

The post-harvest diseases caused to about 50% losses in citrus fruits stored in poor storage conditions, especially under high humidity. The most important fungi causing the post-harvest diseases include: *Penicillium* spp., *Aspergillus* spp., *Alternaria* spp., *Botrytis cinerea*, *Monilinia lax* and *Rhizopus stolonifer* (Agrios, 2005). Most post-harvest pathogens are weak such as (*Penicillium*, *Alternaria*, *Diplodia* and *Phomopsis*) and invade through wounds and when the host defense is weak. Citrus fruits have a pH lower than 4, so most of the fungi attack these fruits (Palou *et al.*, 2001; Milind, 2008 and Zamani *et al.*, 2009). Thermal processing is the most widely used technology for pasteurization of fruit juices and beverages. Juice pasteurization is based on a 5-log reduction of the most resistant microorganisms of public health significance (USFDA 2001

and Rupasinghe, and Juan, 2012). Fruit juice has been traditionally pasteurized by batch heating at 63-65 °C for relatively long time (D'Amico et al. 2006 and Rupasinghe, and Juan, 2012). High temperature short time (HTST) pasteurization is the most commonly used method for heat treatment of fruit juice. Orange juice is processed by HTST at 90 to 95 °C for 15 to 30 s. And apple juice is treated by HTST at 77 to 88 °C for 25 to 30 s (Moyer & Aitken 1980 and Rupasinghe, and Juan, 2012). Sterilizing in an autoclave at 121 °C for 20 minutes (Rupasinghe, and Juan, 2012).

In Egypt and under local markets, there is relatively little information related to the natural occurrence of fungi and mycotoxins in fruits. The investigation aimed to survey the fungal plant diseases in some fruit juices (freshly and pasteurized prepared juices i. e. apple, grape-fruit and navel orange) and their possibility of mycotoxins secretion that may be harmful to humans and its control.

## **Material and method**

### ***Sample collection***

Fresh juice of washington navel orange, (*Citrus sinensis* L.), grapefruits, (*Citrus paradisi* Macf) and apple fruits (*Malus domestica*) as well as pasteurized apple and citrus juices which were used for this work were collected from different supermarket in two different localities i. e. Kalubia and Gharbia Governorates, Egypt. Freshly ripe of apple and citrus fruits (grape-fruit and washington navel orange) were collected in polythene bags and stored at 4 °C till used in the laboratory, Microbiology Department, fac. of Sciences, Zagazige University, for further work.

### ***1-Juice preparation for analysis***

Three juices belong to the commonly citrus fruit consumed such as navel orange, and grapefruits as well as apple fruits were selected for microbiological study. Each sample was washed, peeled, and cut into slices, crushed by a processor to obtain the fruit juices and juice were extracted using the sterile stainless steel blender, then filtered by 1 mm<sup>2</sup> sieve made of stainless steel to obtain 30 L of juice. Concentrated apple and orange juices were diluted with a ratio of 1:1. 2:1 of the juice and poured into sterile beaker then left in room temperature for 5 days to elevate the microorganism count in the specimen according to Torkamani, and Niakousari, (2011) and Kamal *et al.* (2014).

## **2-Microbiological analysis**

Microbial contaminant of apple and citrus juices were done by serial dilution agar plate technique. Ten mL of juice sample was diluted with 90mL of sterilized water (Tournas *et al.*, 2006). One milliliter of fresh and pasteurized of apple and citrus juices sample dilutions was pipetted to Petri dishes containing PDA supplemented with antibiotic (at pH 5.5) for enumeration of fungi in duplicates. These plates were then incubated at  $25 \pm 2$  °C for five days. All tests were performed and the total count experiments were expressed as colony forming units per 1 ml of the sample (CFU/ml) (Torkamani, and Niakousari, 2011 and Iva Dolezalkova *et al.*, 2012). Mould isolates were purified by single spore isolation and maintained on potato dextrose agar, further sub cultured on slants for further study and microscopic examination. Moulds were identified on the basis of morphological and cultural characteristics according to the methods described in “Fungi and Food Spoilage (Kamal *et al.*, 2014 and Vermani *et al.*, 2014) These fungi were observed under a microscope and identification in Plant Pathology Department, National Research Centre, Cairo, Egypt with the help of available literature by Raper, and Fennell, (1965), Ainsworth *et al.* (1972), Biligrani *et al.*, (1991), Barnett and Hunter (1999) Grizzle (2006), Singh (2009). Different frequencies of fungi were noted.

## **3-Mycotoxin production**

The different fungal isolates were propagated as pure culture in 100 ml SMKY broth (Sucrose 200 g, MgSO<sub>4</sub> 7H<sub>2</sub>O 0.5 g, KNO<sub>3</sub> 3 g, yeast extract 7 g) for 10 days. All fungal isolates incubated in dark condition at  $25 \pm 2$  °C.

3.a-The determination of aflatoxins was carried out by using HPLC according to (AOAC, 2007). The HPLC instrument used was waters (474) system, equipped with quaternary pump. The fluorescence detector system was set at 360 nm excitation and 440 nm emission wavelengths. The chromatography column was phenomenex c18 (250x 4.6 mm), 5 µm. The mobile phase system (H<sub>2</sub>O: MeOH: CH<sub>3</sub>CN, 30:60:10 v/v/v) was isocratically at flow rate of 1 ml /min. The data were collected and integrated using Totalchrom Navigator Chromatography Manager Software. according AOAC, (2007); Han *et al.*, (2004) and Embaby *et al.* (2007 and 2012).

3.b-Fusarium mycotoxins assay: The samples were analyzed using the Fumonisin test (FB<sub>1</sub>) by using HPLC, (Ammar, and El-Naggar, 2014).

3.c-Penicillium mycotoxins assay:Patulin in apple juice was tested by Thin Layer Chromatography (TLC) with the advantages of being simple and low cost. One official method of the Association of Official Analytical

Chemists (AOAC) for the analysis of mycotoxins (2007) uses TLC followed by silica gel column chromatography as sample preparation procedures. The analysis was carried out in silica gel plates, using, for detection, the reaction of patulin with 3-methyl-2-hydrazone benzothiazolinone and HCl (MBTH). The limit of detection (LOD) was described as 20 µg/L (Shephard & Leggott, 2000). Alternative detection and quantification methods were described using fluorodensitometry, or absorbance–transmittance followed by densitometry at 275nm. Reversed phase TLC was despite the patulin elution with a wide variety of solvents (Shephard & Leggott, 2000).

Penicillium isolate were examined to produce Patulin. To extract patulin from apples, A small portion of apple was blended for 2 min using a high speed homogenizer in a C<sub>18</sub>-bonded silica solid support and filtered using glass filter paper. Patulin was extracted from homogenized filtrate using acetonitrile: water (5:95 v:v) solution. The solvent was then evaporated at 35 °C under vacuum. The dried residues were dissolved in 1 ml of acetonitrile: water (5:95 v:v) solution according to Begum (2012) and Ammar, and El-Naggar (2014).

#### ***4-Effects of heat treatments on germination (viability) in vitro***

4.a-Hot water treatment: Sterile glass tubes containing 1.8 ml distilled water were placed in water baths at 45, 50, and 55 °C, and allowed to equilibrate for 5 min. afterwards; 0.2 ml of a concentrated spore suspension was added to the tubes, to achieve a final concentration of  $2 \times 10^5$  spore's 1mL. After 5min., tubes were removed from the water baths and placed immediately on ice. Aliquots of the spore suspensions were transferred to Potato Dextrose Agar (PDA), according to Plaza *et al.* (2004), Fatemi and Borji (2011) and Embaby and Hagag (2014). All Petri dishes were incubated for 48 or 72 h at  $25 \pm ^\circ\text{C}$  in darkness.

4.b- Pasteurized apple and citrus fresh juices under agitation to 80 °C De Donno *et al.* (1998) and Sant'Ana *et al.* (2009 and 2010). Orange juice is processed by High temperature short time (HTST) at 90 °C for 30 s. And apple juice is treated by High temperature short time (HTST) at 80 °C for 30 s according to Moyer and Aitken (1980) and Rupasinghe, and Juan (2012) then transferred to Potato Dextrose Agar (PDA). All Petri dishes were incubated for 48 or 72 h at  $25 \pm ^\circ\text{C}$  in darkness.

4.c- Sterilizing apple and citrus fresh juices in an autoclave at 121 °C for 20 minutes then transferred to Potato Dextrose Agar (PDA). All Petri dishes were incubated for 48 or 72 h at  $25 \pm ^\circ\text{C}$  in darkness. The numbers of colony-forming were counted after 3 days of incubation. and the number of viable spores was calculated for each treatment compared with un-treated (as control).

## Results and discussion

### *1-Mycoflora analyses of juices*

**a-Isolation:** Total fungal isolates associated with some Fresh and Pasteurized juices were tabulated in Table (1). Data in this table show that, fungal isolation yielded Three hundred and eighteen fungal colonies were isolated from all the three tested of fresh juices i. e. Apple, Grape fruit and Navel orange and some yeast. Grape-fruit juice was higher total fungal count compare with others, which record 161 fungal colonies equal 50.63% followed by Navel orange juice which gave 126 fungal colonies equal 39.62%. Apple juice was less total fungal count and gave 31 fungal colonies equal 9.75%. On the other hand the same table indicated that, no any fungal colonies were appeared when diluted the same pasteurized juices (Apple, Grape-fruit and Novel-orange) on the same medium. Droby (2006) and Zhu (2006) reported that, plant pathogens may infect fruits either prior to harvest under field conditions or after harvest during transit and storage. It is estimated that about 20-25% of the harvested fruits are decayed by pathogens during post-harvest handling even in developed countries.

**Table 1.** Percentage of total fungal isolates from different juice samples

	Type of juices						Total
	Apple		Grape-fruit		Navel orange		
	F	P	F	P	F	P	
Number of samples	10	10	10	10	10	10	60
Total fungal isolates	31	NF	161	NF	126	NF	318
%	9.75	00.00	50.63	00.00	39.62	00.00	100.00
F = Fresh juice	P = Pasteurized juices		NF=Not found				


**b-Fungal frequency:** Data in Table (2) presented that, seven fungal genera belong to nine fungal species were isolated and identified from the three tested of fresh juices i. e. Apple, grapefruit and navel orange **Fig. (1)**. These are *Alternaria citri*, *Aspergillus flavus*, *A. parasiticus*, *Botryodiplodia theobromae*, *Colletotrichum gloeosporioides*, *Fusarium* sp., *Penicillium expansum*, *P. digitatum*, *P. italicum* and *Rhizopus stolonifer*. Also, data in this table show that *P. italicum* was the most fungal frequency which record 37.42% followed by *P. digitatum* 22.64%, *B. theobromae* 22.01%, *Rhizopus stolonifer* 5.97%, *Alternaria citri* 4.40%, *P. expansum* 3.77%, *Aspergillus flavus* record 1.59%, *A. parasiticus* 1.26% and *Colletotrichum gloeosporioides* 0.63%. Less fungal frequency was recorded with *Fusarium* sp. fungus which gave 0.31%.

Nunes *et al.* (2010) and Munoz *et al.* (2011) reported that, green and blue moulds, due to the pathogenic action of *Penicillium digitatum* and *Penicillium italicum* respectively, are the main causes of citrus fruit losses during postharvest. *Penicillium expansum* causes a soft rot of apples. This particular species has a special significance because of its ability to produce the mycotoxin patulin which has been detected as a contaminant in unfermented apple juices but not in cider. Commercially produced and properly handled bread generally lacks sufficient amounts of moisture to allow for the growth of any microorganisms except moulds. One of the most common is *Rhizopus stolonifer*, of the referred to as the “bread mould”.

*Colletotrichum acutatum* and *C. gloeosporioides* (Penz.) Penz. and Sacc. are the two members of the genus that are most commonly associated with fruit rots in the literature. *Colletotrichum gloeosporioides* is considered a cumulative species and is found on a wide variety of fruits, including apple, almond, avocado, citrus, mango, olive, and strawberry (Arauz, 2000; Embaby and Abd-Ellatife, 2013 and Embaby *et al.*, 2014). The *Colletotrichum acutatum* and *C. gloeosporioides* caused anthracnose in a wide range of hosts found mainly in temperate, subtropical and tropical areas, which causes anthracnose and blight in agriculturally important hosts including many dicotyledonous plants such as apple, strawberry, citrus, and stone fruits (Talhinhas *et al.*, 2005; Lakshmi *et al.*, 2011; Embaby and Abd-Ellatife, 2013 and Embaby *et al.*, 2014).

**Table (2):** Percentage of fungal frequency contaminated some fresh juices

Fungal isolates	Type of juice			Total
	Apple	Grape-fruit	Navel-orange	
<i>Alternaria citri</i>	00	00.00	4.40	<b>4.40</b>
<i>Aspergillus flavus</i>	1.59	00.00	00.00	<b>1.59</b>
<i>A. parasiticus</i>	1.26	00.00	0.00	<b>1.26</b>
<i>B. theobromae</i>	0.63	21.38	0.00	<b>22.01</b>
<i>C. gloeosporioides</i>	0.63	00.00	0.00	<b>0.63</b>
<i>Fusarium</i> sp.	0.31	0.00	0.00	<b>0.31</b>
<i>P. expansum</i>	3.77	0.00	0.00	<b>3.77</b>
<i>P. digitatum</i>	0.00	10.06	12.58	<b>22.64</b>
<i>P. italicum</i>	0.00	16.67	20.75	<b>37.42</b>
<i>R. stolonifer</i>	1.59	2.52	1.89	<b>5.97</b>
Total	9.75	50.63	39.62	<b>100.00</b>



**Figure 1.** See seven fungal genera which were isolated and identified from different fresh juices i. e. *Alternaria* (a), *Aspergillus* (b), *Botryodiplodia* (c), *Colletotrichum* (d), *Fusarium* (e), *Penicillium* (f) and *Rhizopus* (g) respectively

**2-Mycotoxin production:** All the obtained mycoflora i. e. *Alternaria citri*, *Aspergillus flavus*, *A. niger*, *A. parasiticus*, *A. terries*, *Fusarium* sp., *Penicillium degetatum*, *Penicillium italicum* and *Penicillium expansun* were tested for producing mycotoxin(s) by using thin layer chromatography (TLC). Data were recorded in Table (3). Data in this table show that, only one isolate of *Aspergillus flavus* (No. 20) which isolated from apple fresh juice Gharbia sample gave positive producer of Aflatoxin. Also, only one isolate of *Aspergillus parasiticus* (No. 4) isolated from apple fresh juice Gharbia sample gave positive producer of Aflatoxin. Neither *A. niger* nor *A. terries* produced any Aflatoxin and record negative reaction. Kakde and Kakde (2012) reported that, molds, which are of importance in food because of potential mycotoxin production, include members of the genera *Aspergillus*, *Trichothecium*, *Fusarium*. Drusch and Ragab (2003) and Monso (2004) Stated that, some potent fungal toxins like aflatoxins, ochratoxinA, patulin have been detected in fruits during storage.

**Table 3.** Tested of mycotoxin production

Fungal isolate(s)	mycotoxin	juice	Localities	
			Gharbia	Kalubia
<i>A. flavus</i>		apple	+	ND
<i>A. niger</i>	Afs	apple	ND	ND
<i>A. parasiticus</i>		apple	+	ND
<i>A. terries</i>		apple	ND	ND
<i>P. degetatum</i>		citrus	ND	ND
<i>P. italicum</i>	pat	citrus	ND	ND
<i>P. expansun</i>		apple	+	+
<i>Fusarium</i> sp.	F B <sub>1</sub>	apple	ND	ND

Afs = Aflatoxins

F B<sub>1</sub> = Fumonisin B<sub>1</sub>

pat = patulin

ND: Not detected

**Determination of mycotoxin(s)**


Identification and determination of aflatoxions could easily be deduced from the constant retention time compared with the standard spiked in the HPLC chromatogram (Fig. 1). Determination of aflatoxins produced by *Aspergillus flavus* and *A. parasiticus* after inoculated artificially and incubated for two weeks at  $28 \pm 2$  °C were tabulated in Table (4). Data in this table presented that, aflatoxin content after inoculated with the tested of *Aspergillus flavus* isolate No. 20 which isolated from apple fresh juice Gharbia sample was found to be produced only 0.10 ng/ml of aflatoxin AFB<sub>1</sub>(Fig. 2) while, *A. parasiticus* isolate No. 4 isolated from apple fresh juice Gharbia sample was found to be produced all types of aflatoxins which gave 0.52 ng/g of aflatoxin AFB<sub>1</sub>, 0.10 ng/g of aflatoxin AFB<sub>2</sub>, 0.32 ng/ml of aflatoxin AFG<sub>1</sub> and 0.10 ng/ml of aflatoxin AFG<sub>2</sub> (Fig. 3). Pande *et al.* (2012), Sarmah, and Sarma, (2012), Ahire and Sangale (2012) and Vermani *et al.* (2014) reported that, the presence of fungi is a serious health hazard for workers as well as consumers in markets. Quantification of airborne fungi in these markets is essential to identify major pathogenic fungi causing post harvest diseases of fruits. It is crucial for the post-harvest quality management of a wide range of high value fruit crops.

**Table 4.** Quantitively of aflatoxins ng/g HPLC


Juice	Type of		Concentration of aflatoxin (ng/ml)				Total
	Sample	Isolate	AFB <sub>1</sub>	AFB <sub>2</sub>	AFG <sub>1</sub>	AFG <sub>2</sub>	
Gharbia		<i>A. flavus</i>	0.10	ND	ND	ND	0.10
		<i>A. parasiticus</i>	0.52	0.10	0.32	0.10	1.04
Apple	Kalubia	<i>A. flavus</i>	ND	ND	ND	ND	ND
		<i>A. parasiticus</i>	ND	ND	ND	ND	ND
		<i>Fusarium sp.</i>	ND	ND	ND	ND	ND

ND = Not detected


SAMPLE INFORMATION			
Sample Name:	STD	Acquired By:	System
Sample Type:	Unknown	Sample Set Name:	Aflatoxins 2014
Vial:	1	Acq. Method Set:	Aflatoxins 2014
Injection #:	1	Processing Method:	Aflatoxin 2014
Injection Volume:	20.00 ul	Channel Name:	2475ChA ex365/em450
Run Time:	22.0 Minutes	Proc. Chnl. Descr.:	2475ChA ex365/em450
Date Acquired:	10/29/2014 2:17:15 PM EET		
Date Processed:	10/29/2014 2:39:20 PM EET		



**Figure 1.** Standard spiked in the HPLC chromatogram of aflatoxins AF G<sub>1</sub>, B<sub>1</sub>, G<sub>2</sub>& B<sub>2</sub>



**Figure 2.** HPLC chromatogram of aflatoxins produced by *Aspergillus flavus* (No. 20) isolated from apple fresh juice Gharbia sample



**Figure 3.** HPLC chromatogram of aflatoxins produced by *Aspergillus parasiticus* (No.4) isolated from apple fresh juice Gharbia sample

### 3-Effect of heat treatments on spore viability of the tested fungi

a-Effect of different hot water treatments on spore viability of the three tested fungi i. e. *Aspergillus flavus*, *A. parasiticus* and *Pencillium expansum* were recorded in Table (5). Data show that, all hot water treatments were found to reduce the number of spores' viability of the tested fungi. The numbers of spore viability were decreased with increasing the temperature degree used.

Spores viability of *A. flavus* was found to be decreased from 51 to 24 when treated with hot water treatment at 45<sup>0</sup>c with 52.94 %reduction, *A. parasiticus* was found to decrease from 64 to 40 with 37.50% reduction and *P. expansum* from 73 to 6 with 91.78 reduction percent. Hot water treatment at 50<sup>0</sup>c was found to be decreased Spore viability of *A. flavus* from 51 to 13 with 74.51% reduction, *A. parasiticus* from 64 to 40 with 37.50% reduction while, completely decreased was found with *P. expansum* which record hundred percent of reduction. Enhanced effect was recorded when treated Spores viability of *A. flavus* with hot water treatments at 55<sup>0</sup>c which decreased from 51 to 8 with 84.31 %reduction, *A. parasiticus* from 64 to 10 with 84.38 % reduction and *P. expansum* from 73 to zero which gave completely reduction( 100%) .

b-Also, effect of pasteurization and sterilization treatments on spore viability of the tested fungi i. e. *Aspergillus flavus*, *A. parasiticus* and *Pencillium expansum* were recorded in Table (5). Data presented that, both pasteurization and sterilization treatments have a great effect on spore viability of the three tested fungi. Data also show that either pasteurization or sterilization gave completely inhibition of spore viability of the three tested fungi with hundred percent of reduction. In case of *A. flavus* , Pasteurization and sterilization effected on spore viability as spores are completely reduced from (51x10<sup>-4</sup> spores/ml), *A. parasiticus* from (64 x10<sup>-4</sup> spores/ml) and *P. expansum* from (73x10<sup>-4</sup> spores/ml) to zero with 100% reduction.USFDA (2001) and Rupasinghe and Juan (2012) stated that, thermal processing is the most widely used technology for pasteurization of fruit juices and beverages. Juice pasteurization is based on a 5-log reduction of the most resistant microorganisms of public health significance.Fruit juice has been traditionally pasteurized by batch heating at 63-65<sup>0</sup>c (Low temperature long time (LTLT) for relatively long time (D'Amico *et al.*, 2006 and Rupasinghe and Juan, 2012). Pasteurization is the most commonly used method for heat treatment of fruit juice. For example, orange juice is processed by high temperature short time (HTST) at 90 to 95<sup>0</sup>c for 15 to 30 s. And apple juice is treated by HTST at 77 to 88<sup>0</sup>c for 25 to 30 s (Moyer & Aitken 1980 and Rupasinghe, and Juan, 2012) andsterilizing in an autoclave at 121<sup>0</sup>c for 20 minutes (Rupasinghe and Juan, 2012).

**Table 5.** Effect of heat treatments on spore viability of the tested fungi

Tested fungi		Type of treated/ °c									
		Hot water						Pasteurize <sup>d</sup>		Sterilized	
		45 <sup>0</sup> c	%R	50 <sup>0</sup> c	%R	55 <sup>0</sup> c	%R	80 °C	%R	121 °C	%R
<i>A. flavus</i>	51	24	52.94	13	74.51	8	84.31	00	100	00	100
<i>A. parasiticus</i>	64	40	37.50	19	70.31	10	84.38	00	100	00	100
<i>P. expansum</i>	73	6	91.78	00	100.0	00	100.0	00	100	00	100

## Conclusion

Juices squeezed from fresh fruits contain microorganisms which are potentially hazardous to public health. Juices were spoiled with high level of moulds and yeasts which is attributable to low pH of juices. The selling and consumption of juices are never stopped on nutritional grounds as well as livelihood of street vendors. It is alarming situation for suitable agency to take some necessary action, make guidelines to prevent potential food poisoning from juices that contain pathogenic microorganisms and find natural antimicrobials from plants that control spoilage and pathogenic microorganisms in juices. Mycotoxin analysis is an important subject with social relevance. The most suitable analytical methods respecting accuracy, low detection limits and simple procedures for application for the determination of mycotoxins in food products, especially apple juice and its derivatives techniques were thin-layer chromatography (TLC) and high-performance liquid chromatography (HPLC). Heat treatments i. e. waterbath(at 45, 50 and 55 °C), pasteurization (at 80 °C) and sterilization (at 120 °C) treatments were the most suitable for elimination all Juice microbial, this work will contribute to a better characterization and quantification of its presence in human diet.

## References

- Agrios, G. N. (2005). *Plant Pathology*. New York: Academic Press.
- Ahire, Y. R. and Sangale, M. K. (2012). Survey of aeromycoflora present in vegetable and fruit market. *Elixir International Journal* 52:11381-11383.
- Ainsworth, G. C. Sparrow, F. K. and Sussman, A. S. (1972). *The fungi*, 4 A. New York: Academic Press.
- Ammar, M. I. and El-Naggar, M. A. (2014). Screening and characterization of Fungi and their associated mycotoxins in some fruit crops. *International Journal of Advanced Research* 2:1216-1227.
- AOAC (2007). Association of official analytical chemists. official methods of analysis of AOAC. International 17th edition, *Nature Toxins*. Arlington, Virginia, USA: AOAC International.
- Arauz, L. F. (2000). Mango anthracnose: Economic impact and current. *The American Phytopathological Society* 84:600-611.
- Arya, C. and Arya, A. (2007). Aeromycoflora of fruit markets of Baroda, India and associated diseases of certain fruits. *Aerobiologia* 23:283-289.
- Barkai-Golan, R. and Paster, N. (2008). *Mycotoxins in fruits and vegetables*. Academic Press. 395 pp.
- Barnett, H. L. and Hunter, B. B. (1999). *Illustrated Genera of Imperfect Fungi* (fourth edition). Minnesota, USA: APS Press. 218 pp.
- Begum, G. (2012). Ecotoxicology. *InTech*. pp. 75-91.
- Bhale, U. N. (2011). Survey of market storage diseases of some important fruits of Osmannabad District (M.S.) India. *Science Research Reporter* 1:88-91.
- Bilgrami, K. S., Jamaluddin, S. and Rizwi, M. A. (1991). *Fungi of india part III. list and references*. New Delhi: Today and Tomorrow's Printers and Publishers.
- Buck, J. W., Walcott, R. R. and Benchat, L. R. (2003). Recent trends in microbiological safety of fruits and vegetables. *Plant Health Progress*.

- Chen, L., Ingham, B. H. and Ingham, S. C. (2004). Survival of *Penicillium expansum* and Patulin production on stored apple after wash treatments. *Journal of Food Science* 69:669-675.
- D'Amico, D. J., Silk, T. M., Wu, J. and Guo, M. (2006). Inactivation of microorganisms in milk and apple cider treated with ultrasound. *Journal of Food Protection* 69:556-563.
- De Donno, A., Montagna, M. T., Erroi, R., Liaci, D., Sanapo, S. and Caggiano, G. (1998). Food products and fungal contamination. Note II. Study on moulds presence in pasteurized and fresh fruit juices. *Journal of Preventative Medicine and Hygiene* 39:71-73.
- Droby, S. (2006). Improving quality and safety of fresh fruits and vegetables after harvest by the use of biocontrol agents and natural materials. *Acta Horticulturae* 709:45-51.
- Drusch, S. and Ragab, W. (2003). Mycotoxins in fruits, fruit juices, and dried fruits *Journal of Food Protection* 66:1514-1527.
- Embaby, E. M., Abdel-Galil, M. M. and Laila, F. H. (2007). Occurrence of aflatoxins in some rotted apricot fruit in Egypt. *Research Journal of Agriculture and Biological Sciences* 3:631-637.
- Embaby, E. M., Hagagg, L. F. and Abdel-Galil, M. M. (2012). Decay of some fresh and dry fruit quality contaminated by some mold fungi. *Journal of Applied Sciences Research* 8:3083-3091.
- Embaby, E. M. and Abd-Ellatif, A. A. (2013). Species identification of *Colletotrichum* the causal agent of strawberry anthracnose and their effects on fruit quality and yield losses. *Journal of Applied Sciences Research* 9:3845-3858.
- Embaby, E. M. and Hagag, L. F. (2014). Citrus fruit decay and its control. *Journal of Agricultural Technology* 10:1273-1288.
- Embaby, E. M., Layla-Haggage, F., Abdel-MOnem, M. O., Elsayed, T. and Lobna-Al-Awam, R. (2014). Anthracnose disease (*Colletotrichum* sp.) affecting olive fruit quality and its control in Egypt. *Journal of Agricultural Tecnology* 10:1289-1306.

- Fatemi, S. and Borji, H. (2011). The effect of physical treatments on control of *Penicillium digitatum* decay orange cv. Valencia during storage period. African Journal of Agricultural Research 6:5757-5760.
- Grizzle, H. W. (2006). A microtiter plate procedure for evaluating fungal functional diversity on nitrogen substrates. Mycologia 98:353-363.
- Han, D., Macdonald, S. J., Boughtflower, V. and Brereton, P. (2004). Simultaneous determination of aflatoxins and ochratoxin A in food using a fully automated immunoaffinity column clean up and liquid chromatography fluorescence detection. Journal of chromatography A 1059:13-16.
- Herforth, A. W. (2000). An apple a day can functional foods keep the doctor away. Nutrition Reviews 58:1-10.
- Iva, D., Macalik, Z., Butkovicova, A., Janis, R. and Leona, B. (2012). Monoacylglycerols as fruit juices preservatives. Czech Journal of Food Sciences 30:567-572.
- Kakde, U. B. and Kakde, H. U. (2012). Incidence of post-harvest disease and airborne fungal spores in a vegetable market. Acta Botanica Croatica 71:147-157.
- Kamal, R. A., Dhiman, R., Aggarwal, N. K., Kumar, V. and Kaur, M. (2014). Microbes associated with freshly prepared juices of citrus and carrots. International Journal of Food Science 7.
- Lakshmi, B. K. M., Reddy, P. N. and Prasad, R. D. (2011). Cross-infection Potential of *Colletotrichum gloeosporioides* Penz. isolates causing anthracnose in subtropical fruit crops. Tropical Agricultural Research 22:183-193.
- Marchand, L., Murphy, S., Hankin, J., Wilkens, L. and Kolonel, L. (2000). Intake of flavonoids and lung cancer. Journal of the National Cancer Institute 92:154-160.
- Milind, S. L. (2008). Citrus fruit biology, technology and evaluation. First ed book, Copyright © Elsevier Inc. All Rights Reserved Academic Press.
- Monso, E. M. (2004). Occupational asthma in greenhouse workers. Current Opinion in Pulmonary Medicine 10:147-150.

- Moyer, J. C. and Aitken, H. C. (1980). Apple juice. In Nelson, P. E. and Tressler, D. K. (Eds.), Fruit and vegetable juice processing technology. Westport: AVI. pp. 212-267.
- Munoz, A. M., Garcia, B. L., Candelas, A. V. and Marcos, J. F. (2011). Comparative analysis of the sensitivity to distinct antimicrobials among *Penicillium* spp. causing fruit postharvest decay. *Phytopathol Mediterr* 50:392-407.
- Nunes, C., Duarte, A., Manso, T., Garc ía, J., Cayuela, J., Yousfi, K., Mart ínez, M., Weiland, C. and Salazar, M. (2010). Options from integrated management. *Plant Disease* 84:600-611.
- Plaza, P., Usall, J., Torres, R., Abadias, M., Smilanik, J. and Vi ñas, I. (2004). The use of sodium carbonate to improve curing treatments against green and blue moulds on citrus fruits. *Pest Management Science* 60:815-821.
- Palou, L., Smilanik, J., Usual, J. and Vinas, I. (2001). Control of postharvest blue and green moulds of oranges by hot water, sodium carbonate, and sodium bicarbonate. *Plant Disease Journal* 85:371-376.
- Pande, B. N., Dere, P. K. and Arsule, C. S. (2012): Atmospheric fungal diversity over the vegetable market at Aurangabad (M.S.). *Bionano Frontier* pp. 145-150.
- Pitt, J. J. and Hocking, A. D. (1997). Fungi and food spoilage. London: Blackie Academic and Professional. 593 pp.
- Raper, K. B. and Fennell, D. I. (1965). The genus *Aspergillus*. Baltimore: The Williams and Wilkins Co.
- Rathnayaka, R. M. K. (2013). Antibacterial effect of malic acid against *Listeria monocytogenes*, *Salmonella enteritidis* and *Escherichia coli* in Mango, Pineapple and Papaya juices. *American Journal of Food Technology* 8:74-82.
- Rupasinghe, V. H. P. and Juan, Y. L. (2012). Emerging preservation methods for fruit juices and beverages, food additive. Retrieve from: <http://www.intechopen.com/books/food-additive/emerging>.
- Sant'Ana, A. S., Rosenthal, A. and Massaguer, P. R. (2009). Heat resistance and effects of continuous pasteurization on the inactivation of *Byssoschlamys fulva* ascospores in clarified apple juice. *Journal of Applied Microbiology* 107:197-209.

- Sant'Ana, A. S., Simas, R. C., Almeida, C. A. A., Cabral, E. C., Rauber, R. H., Mallmann, C. A., Eberlin, M. N., Rosenthal, A. and Massaguer, P. R. (2010). Influence of package, type of apple juice and temperature on the production of patulin by *Byssochlamys nivea* and *Byssochlamys fulva*. *International Journal of Food Microbiology* 142:156-163.
- Sarmah, P. S. and Sarma, T. C. (2012). Occurrence of aeromycoflora in the fruit markets of Goalpara district (Assam). *The Ecoscan*. 1:299-302.
- Shephard, G. S. and Leggott, N. L. (2000). Chromatographic determination of the mycotoxin patulin in fruit and fruit juices. *Journal of Chromatography A*. 882:17-22.
- Singh, D. and Sharma, R. R. (2007). Postharvest diseases of fruit and vegetables and their management. In Prasad, D. (Ed.), *Sustainable Pest Management*. New Delhi, India: Daya Publishing House.
- Singh, M. P. (2009). Application of biolog FF MicroPlate for substrate utilization and metabolite profiling of closely related fungi. *Journal of Microbiological Methods* 77:102-108.
- Talhinhas, P., Ferreira, P., Neves-Martins, J., Screenivasaprasad, S. and Oliveira, H. (2005). *Colletotrichum acutatum*: principal agente causal da gafa da Oliveira em Portugal. Proceedings of the III Simpósio Nacional de Olivicultura, Castelo-Branco, Portugal.
- Torkamani, A. E. and Niakousari, M. (2011). Impact of UV-C light on orange juice quality and shelf life. *International Food Research Journal* 18:1265-1268.
- Tournas, V. H., Heeres, J. and Burgess, L. (2006). Moulds and yeasts in fruit salads and fruit juices. *Food Microbiology* 23:684-688.
- U.S. Food and Drug Administration (2001). Hazard analysis and critical control point (HACCP); procedures for the safe and sanitary processing and importing of juices; final rule. *Federal Register* 66:6138-6202.
- Vermani, M., Bedi, N. and Hussain, M. S. (2014). Prevalence of culturable airborne fungi in fruit markets of delhi and noida, India. *International Research Journal of Environment Sciences* 3:1-6.

- Zamani, M., Sharifi-Tehrani, A., Ahmadzadeh1, M., Hosseinaveh, V. and Mostofy, Y. (2009). Control of *Penicillium digitatum* on orange fruit combining Pantoea Agglomerans with hot sodium bicarbonate dipping. Plant Pathology 91:437-442.
- Zhu, S. J. (2006). Non-chemical approaches to decay control in postharvest fruit. In Nouredine, B. and Norio, S. (Eds.), Advances in Postharvest Technologies for Horticultural Crops. India: Research Signpost. pp. 297-313.

(Received: 20 January 2014, accepted: 25 February 2015)