
Effect of Weather Parameters on Blue Pine (*Pinus Wallichiana* J.) Needle Blight and Ascospore Release of *Lophodermium Pinastris* in India

Ahanger, F. A. *, Dar, G. H., Beig, M. A., Sofi, T. A. and Ganie, S. A.

Division of Plant Pathology, S. K. University of Agricultural Sciences and Technology of Kashmir, Shalimar, Srinagar, Jammu and Kashmir – 190 025, India.

Ahanger, F. A., Dar, G. H., Beig, M. A., Sofi, T. A. and Ganie, S. A. (2016). Effect of weather parameters on Blue pine (*Pinus wallichiana* J.) needle blight and ascospore release of *Lophodermium pinastri* in India. International Journal of Agricultural Technology 12(6):1099-1112.

Abstract A study was undertaken to assess the effect of meteorological factors on needle blight (*Pinus wallachiana* J.) disease development and ascospore release in Blue pine. The needle blight disease on Blue pine in Kashmir appeared in February and reached maximum by the end of October. During the study period maximum logarithmic infection rate of 0.0618 and 0.0285 unit/day was recorded during the first fortnight of April in 2008 and 2009, respectively. Weather factors *viz.*, temperature and relative humidity were positively correlated with disease intensity (62.1% contribution). However, except temperature, all other weather factors correlated positively with disease development in terms of infection rate (units/day) (25.7% contribution). The correlation studies of various meteorological factors with disease development revealed significant positive correlation of disease intensity with mean temperature, wind speed and relative humidity. Negative correlation of disease intensity with rainfall and wind speed was observed. The apparent infection rate was negatively correlated with temperature, but positively correlated with relative humidity, rainfall and wind speed. The multiple linear regression fitted to data revealed that weather factors influenced the disease intensity to the extent of 62.1% in 2008-2009 and for infection rate to the extent of 25.7%. Correlation of various meteorological factors with spore release revealed that spore release and temperature were negatively correlated during whole season (2009) but positively correlated for August-September period. The correlation of spore release with relative humidity, rainfall and wind speed was significantly positive during whole season. The disease incidence depicted negative correlation with temperature, rainfall and wind speed but was positively correlated with relative humidity. Multiple regression analysis of spores released/cm² and disease incidence with weather parameters revealed the coefficient of determination (R²) of 43.9, 72.6 and 21.0% for the spores released was during June-July, August-September periods and in whole season, respectively. However, it was 35.0, 90.9 and 90.3% for disease incidence during June-July, August-September and in whole season on Blue pine plantations.

Keywords: Blue pine, *Pinus wallichiana*, needle blight, weather parameters

* **Coressponding author** Ahanger, F.A.; **Email:** ahanger123@gmail.com

Introduction

Blue pine or Kail pine (*Pinus wallichiana* Jackson) grows widely in moist and dry temperate forests of Western and Central Himalaya from Jammu & Kashmir to Bhutan at an altitude of 1700 to 3700 m amsl and is one of the main commercial timber tree species grown in Western Himalaya. Blue pine, mainly propagated through nursery-raised seedlings, faces severe regeneration problem under existing agro-climatic conditions. Depending upon the weather conditions, pine trees are often exposed to persistent pathogenic attacks, particularly those causing leaf or needle diseases (Douce *et al.*, 2002). Weather parameters *viz.*, temperature, rainfall, relative humidity, sunshine hours and wind speed play important role in population dynamics and pathogenic distribution so influence the disease incidence. Losses may be severe if weather and field conditions are favourable for disease spread. Since weather influences plant disease epidemics, therefore, understanding of weather data and climatic conditions in relation to disease development is required to have baseline information for developing simple and reliable disease prediction system. Scanty information is available on the influence of environmental conditions conducive for needle blight disease development. Although weather-based disease forewarning models have been developed in various field and horticulture crops, a functionally viable model for disease forecast in forestry crops is needed for effective integrated disease management (Singh *et al.*, 1990, Jayanthi *et al.*, 1993; Prasad *et al.*, 2008). Generally, high humidity ($\geq 90\%$), optimum leaf wetness and conducive temperature ($27 \pm 3^\circ\text{C}$) favour disease development; while wind and rainfall help in spore dispersal to other organs and plants (Picco and Rodolfi, 2002). The impact of environmental conditions and their fluctuations on inoculum build up and disease spread has not been studied quantitatively in needlecast disease. Predictions of environmental factors that have vital role in disease spread are regarded as an economical method for controlling plant diseases.

World-wide the needle blights are caused by several pathogens such as *Scirrhia pini* syn. *Mycosphaerella pini/Dothistroma pini* (red band needle blight); *Scirrhia acicola* syn. *Mycosphaerella dearnessii/Septoria ascicola* (brown spot needle blight); *Lophodermium seditiosum* and *L. pinastri* (Lophodermium needle blight or Lophodermium needlecast); *Ploioderma* spp. (Ploioderma needlecast); *Cercospora pini-densiflora* (Cercospora needle blight); *Cercoseptoria pini* (Cercoseptoria needle blight); *Davisomycellia ampla* (needle blight); *Meloderma desmazierii* (needle blight); *Elytroderma* sp. (Elytroderma needle disease); *Phoma eupyrena* (Phoma blight), etc. (Reddy and Pandey, 1973; Gadgil, 1984; Ivory, 1994; Brown *et al.*, 2003; Jerry and Richard, 2008). Among these, Lophodermium needle blight has posed serious

threat to pine both in nurseries and field (Miller and Watson, 1971), thereby reducing the plant vigour and health by inducing needle discoloration and defoliation (Johansen and Wade, 1987; Ortiz-Garcia *et al.*, 2003; Hanso and Drenkhan, 2008; Kirk *et al.*, 2008). Severe out-breaks of needle blight (*L. pinastri*) have been reported in 1966-67 in several nurseries at Minnesota, Wisconsin and Michigan (USA) which resulted in the destruction of millions of red pine (*Pinus resinosa* Air.) and Scotch pine (*P. sylvestris* L.) seedlings (Skilling and Nicholls, 1974). In Europe, needle blight disease has threatened the nurseries for the last 100 years (Millar and Watson, 1971; Staley, 1975; Minter, 1981). In Himalayan mountainous range the needle blight disease incited by several *Lophodermium* species has severely affected the pine nursery seedlings and plantations in forests, parks and gardens during last few years (Ahanger *et al.*, 2012). Severe infestation in saplings has adversely affected Blue pine regeneration programmes and warrants immediate attention. The information on the meteorological factors affecting the needle blight disease in Blue pine is lacking, especially with respect to Indian subcontinent. The information on the mechanism of disease development is also lacking for proper diagnosis and to develop effective management strategies.

Objectives: The present investigations were aimed at to assess the relationship between various weather factors and needle blight disease in Blue pine and generate baseline information on conidial dissemination which may help in developing appropriate strategies for disease management.

Materials and methods

Effect of meteorological factors on disease

The role of abiotic factors on the progress of needle blight disease in Blue pine was assessed in forest plantations at Forest Nursery, Faculty of Forestry, SKUAST-Kashmir, Shalimar (Srinagar) during the years 2008 and 2009. Six 10-year old Blue pine plants were randomly selected and tagged for scoring of diseased needles under natural epiphytotic conditions. Scoring was done as per 0-4 point scale (Skilling and Nicholls, 1974) wherein 0 = needles/whorls showing no disease symptoms, and 1, 2, 3 and 4 depicted the needles/whorls affected with 1-25, 26-50, 51-75 and $\geq 76\%$ infection, respectively. The scoring was done at fortnightly interval commencing from 2nd fortnight of February during each year and continued till 1st fortnight of November. The per cent disease intensity was calculated as per the formula:

$$\text{Per cent disease intensity} = \frac{\sum (n \times v)}{N \times 4} \times 100$$

where Σ = Summation, n = number of diseased needles/whorls in each

category, v = numerical value of each category, N = number of needles/whorls examined, and 4 = maximum grade value. The growth of disease development in terms of apparent infection rate (unit/day) was calculated as per the formulae given by Vanderplank (1963).

$$r_1 = \frac{2.3026}{t_2 - t_1} \times \log \frac{x_2}{x_1}$$

where, r is logarithmic infection rate (unit/day) and x_1 and x_2 are disease intensity at time t_1 and t_2 , respectively. The data with respect to the weather variables such as temperature, relative humidity, rainfall and wind speed during the study period was procured from the Meteorological Section, Division of Agronomy, SKUAST-Kashmir, Shalimar (Srinagar). The weather data during the study period was correlated with disease development and multiple regressions worked out to determine the influence of individual and combined weather factors on disease development.

Relation between disease incidence, spore discharge and weather parameters

This study was conducted in the year 2009 at Forest Nursery, SKUAST, Shalimar (Srinagar). The study commenced from 2nd June to 30th September, 2009. Twelve 10-year old plantations were randomly selected and on each plantation three vaseline-coated glass slides were hanged on three branches bearing symptomatic needles with hysterothecia. Slides were replaced with new slides at weekly interval and brought to the laboratory to assess the number of spores trapped/cm² of glass slide. The spores trapped were observed under a compound microscope (40x). All the spores, resembling the Description of Minter *et al.* (1982) with respect to *Lophoderemium pinastri*, that appeared in the field during four passes across the width of the slide were tallied. The area observed under each microscopic slide at each magnification was determined and ascospore counts converted to the number of spores/mm². Further, disease incidence was recorded at weekly interval as per the standard method (Skilling and Nicholls, 1974). The meteorological data during the study period was procured from the Meteorological Section, Division of Agronomy, SKUAST, Shalimar, Srinagar (Jammu & Kashmir) and weekly weather data with respect to maximum and minimum temperature, relative humidity, total rainfall and wind speed correlated with the number of spores trapped and percent disease incidence during the corresponding period. Multiple regressions was computed to estimate the effect of combined weather factors on disease development for the whole season as well as at 2 months periodicity taking the number of spores trapped as dependent variable and meteorological factors as independent variables.

Results

Role of weather parameters on disease development

The studies on disease development during 2008 and 2009 and its correlation with weather parameters *viz.*, temperature, relative humidity, rainfall and wind speed revealed that the disease initiated from 2nd fortnight of February and during both the years reached its peak in the month of September (Table 1). In 2008 the disease initiated with intensity of 0.2% in February and attained a peak disease intensity of 25.0% in 1st fortnight of November. No disease progress was noticed beyond this period. The disease trend was almost similar during 2009 with initial disease intensity of 0.86% in 2nd fortnight of February and peak disease intensity of 23.33% in the 1st fortnight of November. During the study period maximum apparent infection rate of 0.0618units/day was recorded during the 1st fortnight of April in 2008 and infection rate of 0.0285 units/day during the same period in 2009.

The mean temperature, relative humidity, wind speed and rainfall varied from 11.68 to 23.90°C, 54.0 to 76.0%, 1.9 to 4.0 kmph and 0 to 4.73 mm, respectively, during infection period in 2008. The temperature, relative humidity, wind speed and rainfall varied from 7.02 to 25.58°C, 56.0 to 71.0%, 1.6 to 3.9 kmph and 0 to 4.94 mm, respectively, during infection period in 2009.

The relationship between weather parameters, apparent infection rate and disease intensity is presented in Table 2. The data revealed that disease intensity significantly and positively correlated with mean temperature ($r = 0.42$) and relative humidity ($r = 0.194$). However, disease intensity exhibited negative and significant correlation with rainfall ($r = -0.15$) and wind speed ($r = -0.79$). In case of apparent infection rate, the temperature (-0.315) and relative humidity (-0.036) depicted a negative correlation. However, rainfall (0.339) and wind speed (0.621) were significantly positively correlated with apparent infection rate.

The association between weather parameters and disease intensity varies with season to season and from year to year so drawing any conclusion would be misleading. Hence we performed regression analysis with pooled data and the conclusions drawn are based only on pooled data analysis. The pooled regression analysis of disease intensity of *Lophodermium pinastri* in relation to weather parameters during 2008-2009 indicated regression coefficients of 0.621. This clearly indicated that 62.1% variation is caused by inexplicable reasons or due to the errors beyond the control of experiments or due to the factors not included in the investigation.

Table 1. Effect of various weather parameters on development of needle blight disease (*Lophodermium pinastri*) in Blue pine during 2008 and 2009

Time of observation		Temperature (°C)	Weather parameters			Pathological parameters	
Month	Fortnight		Relative humidity* (%)	Wind speed* (kmph)	Rainfall* (mm)	Disease intensity (%)	Infection rate (unit/day)
Year 2008							
February	2 nd	5.57	71	2.7	0.89	0.20	0.0098
March	1 st	10.74	60	3.3	0.13	1.36	0.0174
	2 nd	12.01	54	4.0	0.00	2.66	0.0266
April	1 st	11.68	74	2.8	4.73	3.33	0.0618
	2 nd	14.05	57	3.2	2.47	5.86	0.0461
May	1 st	16.65	63	3.2	0.85	7.86	0.0503
	2 nd	17.87	70	3.2	1.75	8.33	0.0461
June	1 st	22.76	68	2.3	1.47	10.33	0.0099
	2 nd	23.17	67	2.4	0.51	12.00	0.0046
July	1 st	22.92	74	2.2	5.21	13.00	0.0165
	2 nd	23.63	65	2.2	0.39	16.66	0.0013
August	1 st	23.90	71	2.4	4.94	17.00	0.0140
	2 nd	22.54	68	2.1	0.35	21.00	0.0030
September	1 st	18.51	70	2.2	2.11	22.00	0.0057
	2 nd	17.41	66	2.3	0.07	24.00	0.0013
October	1 st	15.83	71	2.3	1.06	24.50	0.0012
	2 nd	12.04	76	1.9	1.21	25.00	0.0000
November	1 st	9.84	73	2.17	3.86	25.00	0.0000
Year 2009							
February	2 nd	7.02	67	3.9	1.84	0.86	0.0438
March	1 st	9.84	65	3.7	0.12	1.66	0.0314
	2 nd	9.94	71	3.3	3.09	2.33	0.0194
April	1 st	11.86	71	3.8	4.94	6.66	0.0285
	2 nd	15.38	60	3.8	0.48	7.33	0.0132
May	1 st	15.61	66	3.3	1.77	8.66	0.0193
	2 nd	18.29	60	3.2	1.64	10.00	0.0111
June	1 st	17.14	56	3.7	1.95	12.00	0.0095
	2 nd	20.46	64	2.6	2.51	13.66	0.0205
July	1 st	20.87	64	2.9	0.56	15.00	0.0158
	2 nd	23.76	68	2.4	2.65	17.30	0.0061
August	1 st	25.58	61	2.4	0.53	19.00	0.0045
	2 nd	22.33	64	2.6	1.85	20.33	0.0052
September	1 st	19.69	67	2.4	1.77	22.00	0.0038
	2 nd	20.03	64	2.3	0.03	23.30	0.0001
October	1 st	16.71	67	2.3	0.13	23.33	0.0000
	2 nd	10.81	65	2.3	0.00	23.33	0.0000

November	1 st	11.18	60	1.6	0.00	23.33	0.0000
----------	-----------------	-------	----	-----	------	-------	--------

*Mean of 15 days; Note: During both the years no disease was noticed in 1st fortnight of February

Table 2. Coefficients of simple correlation of meteorological factors with disease intensity and infection rate of needle blight in Blue pine

Weather parameter	Correlation coefficient	
	Disease intensity (%)	Infection rate (unit/ day)
Temperature (°C)	0.420*	-0.315
Relative humidity (%)	-0.194	-0.036
Rainfall (mm)	0.796*	0.621*
Wind speed (kmph)	-0.153	0.339*

*Significant at 5% level

The multiple linear regression equation of pooled data for the years 2008-2009 gave a linear relationship between disease intensity/infection rate (Y) and weather parameters and infection rate (X_1 = temp, X_2 = RH, X_3 = total rainfall, X_4 = wind speed).

$$Y = 49.9 + 0.158 X_1 - 0.149 X_2 - 0.339 X_3 - 10.3 X_4 \dots \dots \dots \text{Eqn.1}$$

$$(R^2 = 0.621; SE = 5.12)$$

$$Y = 0.0415 - 0.000257 X_1 + 0.000201 X_2 + 0.00295 X_3 + 0.0156X_4 \dots \text{Eqn.2}$$

$$(R^2 = 0.423; SE = 0.012)$$

This clearly indicated that a unit change in temperature exerted influence on disease intensity to the extent of 0.158 units in positive direction, followed by relative humidity with -0.149 units, cumulative rainfall with -0.339 units and wind speed with -10.3 units in negative direction.

Role of meteorological factors on disease incidence and spores release

To ascertain the role of meteorological factors in spore release, weather factors such as temperature, relative humidity, rainfall and wind speed were correlated with the number of spores trapped/cm². The study revealed that ascospores were released throughout the season (2nd June to the last week of September, 2009); and the number of spores trapped/cm² varied from 11 to 152 (Fig. 1). Pattern of spore release followed a polynomial curve with a total of three peak spore counts, two in June-July period (9-23 June and 7-28 July), and one in August-September period (1-8 September). Maximum spores release occurred in June-July which was approximately 2.5 times more than August-September period. It seemed that optimum temperature, high relative humidity, wind speed and high rainfall (16.39 mm in comparison to 8.98 mm during August-September) during June-July period favoured maximum spore release (787 spores than 305 spores in August-September). The highest peak of spore release may be attributed to the favourable wet humid weather and favourable

temperature for hysterothecial opening and ascospore release during June-July period.




Figure 1. The number of *Lophodermium pinastri* spores trapped (at weekly interval) and per cent disease intensity (PDI) in relation to various weather parameters during 2009 at Shalimar, Srinagar (J and K)

The correlation between spore release and temperature was significantly negative during June-July period with $r = -0.34$; but was significant and positive in August-September period with $r = 0.17$ (Table 3). However, in whole season the spore release depicted a negative correlation with temperature ($r = -0.30$). The correlation between needle blight incidence and temperature during June-July period was significantly positive ($r = 0.75$) but during August-

Table 3. Coefficients of simple correlation of disease incidence and the number of spores trapped/cm² of glass slide with meteorological factors

Weather parameters	1 st period (June-July)		2 nd period (August-September)		1 st +2 nd period	
	No. of spores trapped/cm ²	Disease incidence (%)	No. of spores trapped/cm ²	Disease incidence (%)	No. of spores trapped/cm ²	Disease incidence (%)
Temperature (°C)	-0.34	0.75*	0.17	-0.79*	-0.30	0.21
Relative humidity (%)	0.23	0.73*	0.47	0.24*	0.06	0.46*
Rainfall (mm)	0.35	-0.61	0.24	-0.10	0.58	-0.67*
Wind speed (kmph)	0.91*	-0.10	0.26	-0.39	0.60*	-0.20

* Significant at 5% level

September period it was significant and negative ($r = -0.79$). However, disease incidence was positively correlated with temperature during the whole season with $r = 0.21$.

The correlation studies between spore release and relative humidity revealed highly significant correlation during August-September period ($r = 0.47$) than during June-July period ($r = 0.23$). The correlation of overall season between spore release and relative humidity was positive with $r = 0.06$. During June-July period the correlation between needle blight incidence and relative humidity was significantly high with correlation coefficient of $r = 0.75$ in comparison to August-September period with $r = 0.24$. The overall correlation between needle blight incidence and relative humidity was significantly positive ($r = 0.46$).

During June-July period the spores release/cm² was found positively correlated with rainfall ($r = 0.35$) followed by August-September period ($r = 0.24$). The results indicate that spore release over a season was highly significant with rainfall with a correlation coefficient of ($r = 0.59$). Needle blight incidence showed negative relationship with rainfall throughout the season with correlation coefficients of -0.61 , -0.10 and -0.67 during June-July, August-September and overall season, respectively.

The correlation between spore release and wind speed was positive and highly correlated ($r = 0.91$) during June-July period followed by whole season ($r = 0.60$) than in August-September period ($r = 0.26$). However, disease incidence depicted negative correlation with wind speed with correlation coefficients of -0.10 , -0.39 and 0.20 in June-July, August-September and overall season, respectively.

Table 4. Multiple regression equations indicating the relationship of meteorological factors with disease incidence and the number of spores trapped/cm² glass slide

Period of observation	Linear equations	Coefficient of determination {R ² } (%)
Season	ST = 375 + 2.61 T + 3.06 H + 8.89 R + 66.1 W	43.9
	PDI = 35.9 - 0.40 T + 0.15 H + 0.08 R - 8.21 W	35.0
June-July	ST = 125 + 2.39 T + 1.30 H + 15.8 R + 23.9 W	72.6
	PDI = 45.7 + 1.04 T + 0.40 H - 0.07 R + 3.75 W	90.9
August-September	ST = 324 + 2.64 T + 3.49 H - 1.11 R - 32.1W	21.0
	PDI = 14.8 - 0.81 T + 0.18 H - 1.48 RF + 5.45 W	90.3

Note: ST = Number of spores trapped/cm² glass slide, T = mean temperature, H = mean relative humidity (%), W = Mean wind speed, R = mean rainfall, PDI = Percent disease incidence; and R² = regression coefficient.

Multiple regression analysis of the number of spores released/cm² and disease incidence with the prevailing weather parameters was analyzed to know

the extent of correlation in June-July, August-September and over the whole season during 2009. The multiple linear regression of spore release/cm² in relation to weather parameters during 2009 indicated that regression coefficients for mean temperature, relative humidity, rainfall and wind speed in whole season were 2.61, 3.06, 8.89 and 66.10, respectively (Table 4). During June-July the regression coefficients for these weather parameters were of 2.39, 1.30, 15.80 and 23.90, respectively. The regression coefficients during August-September period for mean temperature, relative humidity, rainfall and wind speed were 2.64, 3.49, -1.11 and -32.10, respectively. The coefficient of determination (R²) for the number of spores released was 43.9, 72.6 and 21.0% during June-July, August-September and in whole season, respectively.

The multiple linear regression of Lophodermium blight incidence of *Pinus wallichiana* in relation to weather parameters during 2009 indicated regression coefficients of -0.40, 0.15, 0.08 and 8.21 for mean temperature, relative humidity, rainfall and wind speed, respectively, in the whole season. The regression coefficients during June-July for mean temperature, relative humidity, rainfall and wind speed were 1.04, 0.40, -0.07 and 3.75, respectively, and the regression coefficients for these weather parameters during August-September were -0.81, 0.18, -1.48 and 5.45, respectively. The coefficient of determination (R²) for needle blight incidence on Blue pine plantations during June-July, August-September and in whole season were 35.0, 90.9 and 90.3%, respectively.

Discussion

Precise studies on climatic requirements of Lophodermium needle blight disease is lacking and the knowledge about climatic requirements may help in effective management of disease through timely adaption of control strategies. The study clearly indicated that temperature, relative humidity, wind speed and rainfall influence disease expression; and favourable temperature and wind speed are important weather parameters for the development of needle blight disease. The maximum disease development during early summer could be attributed to the favourable temperature coupled with higher relative humidity and frequent rains which coincided with the formation of primary needles of Blue pine. Warm moist summers and cool wet falls and abundance of young susceptible hosts reportedly favour disease development (Woods, 2003). The findings are supported by Hanso (1970); Pagony (1971), Raspopov (1980); Aminev (1980) and Hanso and Hanso (2001) who found warm wet weather in early spring conducive for disease development on pine plantations. Pagony and Gasko (1977) also observed epidemics of disease due to high seedlings

density and moisture. Multiple regression analysis showed that the weather factors accounted for 62.1% variation in disease development, whereas infection rate accounted for 42.3% disease development. The results are in agreement with Kulkarni (2009) who observed similar type of regression equations for anthracnose disease of green gram and reported that weather factors influence the disease to the extent of 57%.

To ascertain the role of meteorological factors in spore release, weather factors such as temperature, relative humidity, rainfall and wind speed were correlated with the number of spores trapped/cm². The study revealed that the ascospores were released throughout the season showing maximum spores release in June-July than in August-September period. It seemed that optimum temperature, high relative humidity, wind speed and high rainfall during June-July period favoured maximum spore release (787 spores than 305 spores in August-September). The highest peak of spore release may be attributed to the favourable wet humid weather and favourable temperature for hysterothecial opening and ascospore release during June-July period. Maximum spore release seemed to be related with the prevailing temperature conditions after the periods of maximum rainfall. However, during August-September period though prevailing temperature were favourable but rainfall was less which delayed ascospore release of *L. pinastri* so resulted in less disease during the period. These findings are in agreement with Chwalinski (1978), Fedorov and Yakimov (1979) and Raspopov (1980) who observed ascospore release throughout the season (from April to November) and found two spore peaks one in summer (May-June) and another in autumn (August to mid-November). Pagony and Gasko (1977) also reported that the number of spores collected in spring-summer period were 5 times higher than those in summer-autumn. Minter and Miller (1980) reported that hysterothecia open during wet periods and release ascospores. They observed earlier spore dispersal in *L. pinastri* and *L. conigenum* for one to few months than observed in other related species. Raspopov (1980) reported that ascospore dissemination dynamics depends on the number of apothecia with mature ascospores, air and temperature and needle humidity and observed increase in ascospore release at 15-20°C and decreases at 10-12°C. He also recorded significant increase in spore dissemination after daily precipitation of 8-10 mm. Linzon (1968) have also reported highest number of ascospore release during the month of June compared to July-August as result of maximum rainfall during June.

The correlation between spore release and temperature was significantly positive in whole season and in August-September period but negatively correlated in June-July period. Spore release was highly correlated with relative humidity in August-September period than during June-July period. During

June-July period the spore release/cm² was positively correlated with rainfall followed by August-September period. The results indicated that spore release over a season was highly significant with rainfall. Needle blight incidence showed negative relationship with rainfall throughout the season during June-July, August-September and overall season, respectively. The observations are in corroboration with Ostry and Nicholls (1992) who reported positive correlation between rainfall and spore release of *L. pinastri*. Similarly, Chwalinski (1978) reported maximum ascospore production and dispersal as well as highest infection rate in autumn season in case of *L. pinastri*. Pagony and Gasko (1977) observed maximum spore discharges in mid June to mid July and mid August to ending September. Further, the intensity of spore dissemination was higher in young than in mature pine plantations. Pagony (1971) observed a positive correlation between rainfall and the number of spores trapped during the growing period. Scharpf (1986) reported a maximum ascospore release (1,170 spores) on 10 spore traps in the 1st week of May, 1988 during which 5 cm rainfall was recorded.

Multiple regression analysis of the number of spores released/cm² and disease incidence with the prevailing weather parameters was analyzed to know the extent of correlation in June-July, August-September and over the whole season during 2009. The multiple linear regression of spore release/cm² in relation to weather parameters during 2009 indicated that regression coefficients for mean temperature, relative humidity, rainfall and wind speed in whole season were 2.61, 3.06, 8.89 and 66.10, respectively. During June-July the regression coefficients for these weather parameters were of 2.39, 1.30, 15.80 and 23.90, respectively. The regression coefficients during August-September period for mean temperature, relative humidity, rainfall and wind speed were 2.64, 3.49, -1.11 and -32.10, respectively. The coefficient of determination (R²) for the number of spores released was 43.9, 72.6 and 21.0% during June-July, August-September and in whole season, respectively. The coefficient of determination (R²) for needle blight incidence on Blue pine plantations during June-July, August-September and in whole season were 35.0, 90.9 and 90.3%, respectively. The results are in agreement with Kulkarni (2009) who observed similar type of regression equations for anthracnose disease of green gram and reported that weather factors influence the disease to the extent of 57%. Aminev (1980) developed a forecast system for *L. pinastri* on the basis of correlation between the development of needle blight disease in a year and the sum of monthly rainfall, snow depth and available monthly temperature from primary infection period to its renewal in the following year.

- Nicholls, T. H. and Skilling, D. D. (1974). Control of *Lophodermium* needlecast disease in nurseries and Christmas tree plantations. USDA Forest Service Research Paper, NC-110. 11 pp.
- Ortiz-García, S., Gernandt, D. S., Stone, J. K., Johnston, P. R., Chapela, I. H., Salas, L. R. and Alvarez, B. (2003). Phylogenetics of *Lophodermium* from pine. *Mycologia* 95:846-859.
- Ostry, M. E. and Nicholls, T. H. (1992). A technique for trapping fungal spores. USDA North Central Forest Experimental Station, St Paul, Minnesota Research Note. NC-283, pp. 1-4.
- Pagony, H. and Gasko, E. (1977). Controlling effect of Scotch pine by *Lophodermium pinastri* in nurseries. *Erdo* 26:425-429.
- Pagony, H. (1971). The course of the weather and control of the *Lophodermium* needlecast. *Przebieg Pogody a Zwalczenie Osutki Sosny* 127:109-116.
- Raspopov, P. M. (1980). Dates of ascospore discharge of the pathogen of needlecast in the south Ural and optimal times for prophylactic sprayings. *Mikologiya i Fitopatologiya* 14:440-445.
- Scharpf, R. F. (1986). Effect of a foliage disease caused by *Lirula abietisconcoloris* on growth of white fir in California. *Plant Disease* 70:13-14.
- Skilling, D. D. and Nicholls, T. H. (1974). Brown spot needle diseases: biology and control in Scotch pine plantations. USDA Forest Service Research Paper NC-109.
- Staley, J. M. (1975). The taxonomy of *Lophodermia* on pines with special reference to problems in North American Christmas tree plantations. *Mitt. Bundesforschungsanst. Forst-Holzwirtschaft* 108:79-85.
- Vanderplank, J. E. (1963). *Plant Disease: epidemics and control*. Academic Press, New York, USA. 349 pp.
- Woods, A. J. (2003). Species diversity and forest health in Northwest British Columbia. *The Forestry Chronicle* 79:892-897.

(Received: 8 July 2016, accepted: 5 November 2016)