
Effect of Seed Priming with Salicylic Acid on Seed Germination and Seedling Growth of Broad bean (*Vicia faba* L).

Soliman, M. H. *, Al-Juhani, R. S., Hashash, M. A. and Al-Juhani, F. M.

Biology Department, Faculty of Science Yanbu, Tabiah University, Saudi Arabia.

Soliman, M. H., Al-Juhani, R. S., Hashash, M. A. and Al-Juhani, F. M. (2016). Effect of seed priming with salicylic acid on seed germination and seedling growth of broad bean (*Vicia faba* L). International Journal of Agricultural Technology 12(6):1125-1138.

Abstract Seed priming with growth regulators, especially salicylic acid, is a promising alternative seed primer because it is an important inducer of resistance against diseases and pests, as well as acting significantly on quality and seed yield. The exogenous salicylic acid SA applied at 0.0, 0.5, 1.0, and 3.0mM SA concentrations to faba bean seeds showed significant increase in seed germination and seedling growth except at high concentrations. Seed germination decreased by increasing of SA up to 3.0mM. As, SA at high concentrations 1.0 and 3.0 mM had inhibitory effects on germination as well as it reduced seedling growth, where the lengths of plumule and root (radical) were reduced compared to control. Moreover, these highest concentrations of SA shown negative result for fresh root weight. Meanwhile, SA at low concentrations had no significant effect at ($P>0.05$) on all tested growth parameters. It can be noted seed priming of faba bean with low concentration of SA will speed up the germination time and enhance the establishment of seedlings as well as powerful tool in enhancing the growth and productivity of such crop especially in areas with saline irrigation water resources. Moreover; primed seeds will be less susceptible to soil-borne pathogens.

Keywords: salicylic acid, germination, seed vigor, growth regulator

Introduction

Faba bean (*Vicia faba* L.) is beneficial for human nutrition perspectives because they are good source of energy, protein and dietary fiber Ma *et al.*, (2005). Seed vigour is a very good indicator of the potential field performance followed by the field planting value. Growth regulators play an important role in increasing germination rate of seeds, employed for rooting, vegetative propagation and overall yield of several plants as well as they mostly used for improving seed germination and the productivity of a large number of agricultural crops Jam *et al.* (2012).

Salicylic acid SA is a natural compound that plays a central role in certain physiological processes and defense responses in plants Shi and Zhu, (2008). It can affect seed germination, cell growth, stomatal opening, expression of genes associated with senescence, and fruit production

Corresponding author: Soliman, M. H.; **Email:** monash12@gmail.com

Klessig *et al.* (2009). Salicylic acid plays an important role for increasing resistance to environmental stress Raskin (1992). It has been reported that SA increases salinity tolerance Jam *et al.* (2012) and resistance to water deficit in seedlings Bezaukova *et al.* (2001). The use of these regulators increases plant tolerances to abiotic and biotic stresses.

Many researchers have evaluated the processes involved in seed germination, and how they are affected by plant hormones in a range of plant families Hermann *et al.* (2007). In drought condition, using some plant growth regulators such as salicylic acid and cycocel might be an effective strategy to prevent destructive effect of drought and provide plant compatibility. These regulators are known as a strategy to prevent the harmful effect of environmental stresses. These stresses include: heat, cold, heavy metals and drought. Salicylic acid is part of a signaling pathway that is induced by a number of biotic and abiotic stresses. A lot of data exist on the protective effect of SA against ultraviolet light Yalpani *et al.* (1994), salinity Shakirova *et al.* (2003), drought Singh and Usha (2003), heavy metal toxicity Metawally *et al.* (2003) and mediating plant defense against pathogens Raskin (1992).

Materials and methods

The experiments were carried out under laboratory conditions at Department of Biology, Faculty of Science, Yanbu Branch, Girls Sections, Taibah University, Kingdom of Saudi Arabia during February and March 2016.

Source of Seed Samples and Chemical Substance (SA)

Seeds of Faba bean (*V. faba* L.) were purchased from a local supermarket in Yanbu, Kingdom of Saudi Arabia. Broad bean seeds were first surface sterilized with 0.2% mercuric acid for 10 min and followed by washed repeatedly with sterile water to remove all traces of HgCl₂. Salicylic acid (Sigma, Aldrich) is obtained from Laboratory of the Department of Chemistry, Faculty of Science, Yanbu Branch, Girls Sections, Taibah University, Kingdom of Saudi Arabia .

Germination Bioassay

The seeds of faba bean were soaked in distilled water to test their viability, and then the precipitated seeds were air dried at room temperature. Healthy seeds of the recipient species were disinfected with 0.2% mercuric acid for 10 min and followed by washed repeatedly with sterile water to remove all traces of HgCl₂.

Different concentrations of exogenously applied Salicylic acid SA i.e., 0.5 mM, 1.0 mM and 3.0 mM) are used as a substrate medium for the germination experiment. Fifteen healthy seeds (in three replicates) for the recipient species during the experimental period were allowed to germinate in the different concentrations of SA under normal laboratory conditions with day temperature ranging from 20-22°C and night temperature from 14-16°C. Sterilized petri-dishes (150×20 mm diameter) were used for germination test. Fifteen seeds of nearly equal size were placed in each Petri dishes double laid with Whatman No1 filter paper. Fifteen to twenty ml of each level of the donor treatment solutions were added to each of the petri-dishes respectively for different concentration. Distilled water was used as a control.

Petri dishes were incubated in a lit room at an average temperature of about 20-22°C for 9 days. The experiment was replicated 3 times. Germination inhibition or stimulation percentage was calculated for each concentration treatment according to the general equation. Germination percentage was determined according to recommended methods by (ISTA, 1985). Number of germinated seeds was counted starting from the 1st day after germination when roots appeared till the 9th day, which is the last day. Emergence of radical is considered as an indicator of germination. *Germination Percentage was calculated as follows* $GP = \frac{\text{Number of germinated seeds}}{\text{Number of total seeds}} \times 100$.

Seed Viability and Seedling Vigour index

Seed germination tests were conducted for each solution of Salicylic acid SA as maintained above as follows: Standard, Root length, plumule length and Vigour index determined following method of Abdul Baki and Anderson (1973). Seedling vigour was estimated by measurement of the length of germinated roots and shoots daily from the 3rd day after planting till the 9th day, then adding the recorded measurement taken during the experimental time and multiplying with germination percentage. Seedling vigour index was determined as the product of the percentage germination and that of seedling length. *Seedling vigor index (SVI) was calculated by the following formula* (Abdul-Baki and Anderson, 1973):

$$SVI = \frac{\text{Seedling length cm} \times \text{Germination percentage}}{100}$$

Determination of Fresh matter

Remove seedlings from Petri-dishes and wash off any filter paper remains. Blot plants gently with soft paper towel to remove any free surface moisture. Weigh immediately (plants have a high composition of water, so waiting to weigh them may lead to some drying and therefore produce inaccurate data).

Statistical analysis

Data concerning the effect of different concentrations of salicylic acid on faba bean_ seed germination and seedling growth was subjected to standard analysis of variance (ANOVA) using SPSS V.16 and with the least significant difference was used to compare means of traits ($p < 0.05$).

Results

Seed Viability and Vigor Index Test

Different concentrations of Salicylic acid SA were applied as seed primer of faba bean. Results in Table 1 showed that the Effect of the four SA concentrations on vigour index of faba bean at the end of priming period (9 days). The results indicate that all tested concentrations of SA (0.5 mM , 1.0 mM and 3.0 mM) affected vigour index of faba bean. 0.5mM and 1.0 mM of SA the cause highest records of vigour index of faba bean .

Meanwhile, high concentration of SA (3.0 mM) cause a significance decrease in vigor index when compared with control treatments. As concentration of SA increased the vigour index significantly decreased. Maximum vigour index recorded in 0.5mM SA compared to control sets (2277 ± 4.198); whereas the maximum inhibitory effect of the donor treatment (SA) for seed vigor index was recorded in 3.0 mM SA (309 ± 59.91).

Table 1. Effect of Different Concentrations of Salicylic acid SA on the Seed Vigor Index (SVI) of Germinated faba bean_Seeds

Salicylic acid SA Concentration	Seed Vigor Index /Days				
	1 st day	3 rd day	5 th day	7 th day	9 th day
Control	0	86 a	280 a	839.2 b	1520 c
0.5 mM	0	108 b	533 b	119.5 a	2277 b
1.0 mM	0	63.84 a	240 a	596.5 b	708.3 a
3.0 mM	0	40 a	114.4 a	253.1 a	309.7 a

Figures with the same letter are not significantly different ($p=0.05$)

Seed germination

Effect of different concentrations of SA on germination and inhibition percentage (GP or IP) of faba bean was represented in (Table 2). The results indicate that maximum percentage of germination was reported in the concentration 0.5mM of SA ,which was (93.3±16.13145%) compared to that of all the concentrations and control treatments which was reported (80 ±14.661%).

Table 2. Effect of Different Concentrations of Salicylic acid SA on the Percentage of Germination (GP) of faba bean Seeds.

Salicylic acid SA Concentration	Germination Percentage (GP) /Days				
	1 st day	3 rd day	5 th day	7 th day	9 th day
Control	0	20 a	40 b	66.6 c	80 c
0.5 mM	0	33.3 b	53.3 b	73.3 c	93.3 c
1.0 mM	0	26.6 b	40 b	46.6 b	46.6 b
3.0 mM	0	20 a	26.6 a	33.3 a	33.3 a

Figures with the same letter are not significantly different (p=0.05)

Whereas; the minimum percentage of germination (Maximum inhibition) of seed germination was found on higher concentration of SA showing only 33.3±6.17451% germination.

Root Length (RL)

Compared to the control, a gradual decrease in radicle (RL) lengths of faba bean seedling roots was noticed along the gradual SA concentrations (Table 3). Obviously, all concentration levels had significantly decrease of faba bean seedling roots RL The maximum root length recorded for 0.5mM SA was (13.1± 2.35 cm) compared to control which (10± 1.757cm) and it was followed by 1mM SA as (7.2± 1.54cm) respectively while a minimum root length was found at highest SA concentration 3mM (4.8± 1.81cm). As the concentration of SA decreased, there was an increment in root length.

Table 3. Effect of Different Concentrations of Salicylic acid SA on Root Lengths (RL) (cm) of faba bean Seedlings

Salicylic acid SA Concentration	Root Length (cm) /Days				
	1 st day	3 rd day	5 th day	7 th day	9 th day
Control	0	1.8 b	3.0 a	6.0 b	10.0 b
0.5 mM	0	2.0 b	6.0 b	8.0 b	13.1 b
1.0 mM	0	1.4 a	4.0a	7.5bc	7.2 a
3.0 mM	0	1.2 a	3 a	4.3 a	4.8 a

Figures with the same letter are not significantly different (p=0.05).

Plumle Length (PL)

Results in (Table 4) showed that primed faba bean seeds with SA increased the plumule length of faba bean seeds compared with high concentrations and control treatments control. Whereas, the minimum plumule length was found at highest concentration (3.0 mM SA) which was (4.5± 0.238cm) while the maximum was given by 0.5 mM SA (11.3± 2.13cm). As the concentration of SA decreased, there was an increment in plumule length.

Table 4. Effect of Different Concentrations of Salicylic acid SA on Plumle Lengths (PL) (cm) of faba bean Seedlings

Salicylic acid SA Concentration	Plumle Length (cm) /Days				
	1 st day	3 rd day	5 th day	7 th day	9 th day
Control	0	2.5 c	4.0 c	6.6 b	9.0 c
0.5mM	0	1.2 a	4.0 c	8.3 c	11.3 c
1.0 mM	0	1.0 a	2.0 b	5.3 b	8.0 b
3.0 mM	0	0.8 a	1.3 a	3.3 a	4.5 a

Figures with the same letter are not significantly different (p=0.05).

Root Fresh Weight (gm)

Seedling fresh weight of broad bean are showed in (Table5). Data reveals that; the inhibitory effect was concentration dependent, as a gradual inhibition in root fresh weight obtained as concentration increased. Results of root fresh weight reveals that priming *faba bean* seeds with SA decreased the root weight as compared to control, Whereas; at the end of priming period (9 days) the highest root weight (11.3 ± 2.133 gm) was recorded by 0.5mM SA in comparison with control priming (9 ± 1.567 mg) and it was followed by 1mM SA, and 3mM SA (8 ± 1.482 gm, 4.5 ± 0.768 gm) respectively. On the other hand, minimum root fresh weight was given by 3mM SA (4.5 ± 0.768 gm).

Table 5. Effect of Different Concentrations of Salicylic acid SA on Root Fresh Weight (gm) of faba bean Seedlings

Salicylic acid SA Concentration	Root Fresh Weight (gm) /Days				
	1 st day	3 rd day	5 th day	7 th day	9 th day
Control	0	2.5 c	4.0 b	6.6 c	9.0 c
0.5 mM	0	1.2 a	4.0 b	8.3 d	11.3 d
1.0 mM	0	1.0 a	2.0 a	5.3 b	8.0 b
3.0 mM	0	0.8 a	1.3 a	3.3 a	4.5a

Figures with the same letter are not significantly different ($p=0.05$).

In conclusion, According to the results of this study and the ANOVA analysis it could be noted that, no statistically significant between 0.5 and 1.0 mM of SA concentrations at the 9th day after sowing. Based on the results, it can be concluded that low concentrations of SA have more positive effects on faba bean germination than medium and high concentrations. This means that seed priming of faba bean with this low concentration of SA will speed up the germination time and enhance the establishment of seedlings.

Discussion

The results of this study clearly show that, the effect of salicylic acid on the seed germination and seedling growth of faba bean was stated to be variable, promoting some parameters and inhibiting others depending on its

concentration. As, Maximum vigour index recorded in 0.5mM SA compared to control. Whereas the maximum inhibitory effect of (SA) for seed vigor index was recorded in 3.0 mM . In our treatments with increased SA concentration the percentage of germination and seedling growth parameters in the different treatments during 9 days of faba bean seeds were decreased. On the contrary, SA enhanced the emergence in lower concentrations; that appeared harmless for both seed germination and establishment of the seedling.

The promotive effect of salicylic acid could be attributed to its bioregulator effects on physiological and biochemical processes in plants such as:- Promoting effect of SA is due to increased level of Cell division, cell differentiation and cell elongation within the apical meristem of seedling root, and increment in biosynthesis of organic foods which caused an increase in seedling growth (Abd El-Wahed *et al.*, 2006; El-Khallal *et al.*, 2009, and Delavari *et al.*, 2010). Ion uptake, transport and the availability and movement of nutrients could result in stimulating different nutrients in the leaves (Raskin, 1992). Habibi and Abdoli (2013) reported that low concentrations of SA increased the germination percentage, but this increase was not significant compared to the control treatment. Similar, Farahbakhsh (2012) reported that the concentration of 0.25 and 0.5 mM of salicylic acid on measured traits was more effective compared with the other levels. There are conflicting reports about the effects of salicylic acid on seed germination. Sharafizad *et al.* (2013) reported that after the control treatment (priming with distilled water), the highest percentage of germination was for low concentrations of Salicylic acid (0.7 mM) and germination percentage decreased with increasing concentrations of SA. These in agreement with our results. SA induced several effects depending on the concentration applied and quite high doses were required to observe symptoms; in all cases, the first effect observed was a delay in seed germination (Metraux *et al.*, 1990). Above a concentration of 0.5mM, the seed germination process was dramatically impeded.

Our results also indicate that, the minimum percentage of germination (Maximum inhibition) of seed germination was found on higher concentration of SA. In this respect, Several workers reported that stimulating effects of SA on germination are concentration dependent (Rajjou *et al.*, 2006 and Singh *et al.*, 2010). SA significantly stimulated the activities of enzymes involved in germination such as transketolase, enolase, malate dehydrogenase, phosphoglycerate kinase, glyceraldehyde 3-phosphate, dehydrogenase, fructose 1,6-diphosphatase, and pyruvate decarboxylase. In addition; seeds germinated in SA as substrate media showed abundant levels of isocitrate lyase and malate synthase (key enzymes of glyoxylate cycle) (Eastmond and Graham, 2001 and Rajjou *et al.*, 2006). Rajjou *et al.* (2006) hypothesized that detoxification mechanism in germinating seeds counteract by exogenous SA treatments.

SA significantly increases plant growth either in stress or without stress conditions. This effect probably related to SA inhibition of Cl⁻ and Na⁺ absorption and its help for Mg, Fe, Mn, N and Cu absorption or results from its effect on lipid peroxidation and membrane permeability (Gunes *et al.*, 2007). Lower SA could promote sink/source regulation, enzymatic activities, protein synthesis and photosynthetic activity as well as increase the antioxidant capacity of plants (Raskin, 1992; Blokhina *et al.*, 2003 and El-Tayeb, 2005). It is possible that SA stimulates the seed germination via bio-synthesis of gibberellic acid and acts as thermogene inducers Shah, (2003). Eastmond and Graham, (2001) reported that isocitrate lyase and malate synthase were found to be more abundant in seeds germinated in the presence of SA than in its absence. Isocitrate lyase and malate synthase are two key enzymes of the glyoxylate cycle that play a crucial role in the synthesis of carbohydrates from storage lipids during seed germination and seedling establishment. An elicitation of these metabolic pathways by SA would favor the transition from a metabolically quiescent state to a metabolically active state during germination, thereby accounting for the observed improvement in seed vigor in the presence of this molecule. SA caused marked increments in IAA, GA3, zeatin and zeatin riboside, in the meantime decrease in ABA content comparing with untreated controls (Shehata *et al.* (2001) and Zaghlool (2002). On the other hand, the decrease in ABA content attributed to the shift of the common precursor isopentenyl pyrophosphate to biosynthesis of cytokinins and/or gibberellins instead of ABA (Hopkins and Huner, 2004). Findings of Eisavand *et al.* (2011) showed an increase in respiration activities, ATP production, induced RNA activity and protein synthesis in primed seeds, as well as enhanced radicle length and weight of primed sunflower seed.

Results in this study are in agreement with several reports which related to applications of SA at higher levels could reduce emergence index. Similar results have been reported for maize embryo germination, for which high doses of SA, in the range of 3 to 5 mM, completely inhibited germination (Guan *et al.*, 2007). Above a concentration of 0.5 mM, this compound entailed a strong retardation of growth and plants appeared bleached, most presumably because at these high concentrations SA induced an oxidative stress (Rao *et al.*, 1997). Improvement in germination characteristics of primed seeds could be results of increasing the antioxidant profile of treated seeds (Ansari and Sharif-Zadeh, 2012). Increasing SA concentration caused many changes in physiological pathways during seed germination and seedling growth. From these changes:-When exogenous SA is more than 1 mM, plants usually trend to oxidative burst and cell death. On the basis of evidence that high levels of SA can bind and inhibit H₂O₂ removing enzymes, ascorbate peroxidase (APX) and catalase (CAT) (Durner and Klessig, 1995). Findings of Nun *et al.* (2003) showed that SA can inhibit the activity of catalase. Reduced catalase activity leads to

increased hydrogen peroxide (H_2O_2) product which stops germination processes. At the higher levels of SA; sucrose which come from scutellum independently to seedlings during germination is inhibited. As a result ketoaldehydes, H_2O_2 and various free radicals are generated and inhibit seedling growth (Russell *et al.*, 2002). Enhance ABA synthesis, which can stop the seed germination (Wu *et al.*, 1998; Ahmad *et al.*, 2012). In addition, NO (Nitric Oxides) can chemically react with superoxide, generating the cytotoxic compounds peroxynitrite and hydroxyl radical, therefore inducing oxidative burst and cell death (Van Camp, *et al.*, 1998). SA is an effective inhibitor of ethylene biosynthesis and does it via effect on ACC- synthetase enzyme (Raskin *et al.*, 1992). Results represented in this research; the decrease in germination percentage may be result of ethylene biosynthesis inhibition. Mazaherie Tirani and Manouchehri (2005) reported that high concentrations of SA reduced Brassica napus seed germination compared to hydro-priming which is in accordance with the findings of the present study. Whereas, When using 0.5 mM salicylic acid, Szepesi *et al.* (2005) found increased percentage of germination in tomato seeds. Data agree with those found by Maia *et al.* (2000), who have found in soybean that salicylic acid used alone has increased the lengths of shoots and roots and fresh weight at concentrations of 50 and 100 mg.kg⁻¹. Reduction in physiological parameters previously reported in our study may be a result of higher dosages of SA used where toxic effects are more pronounced than beneficial effects of SA and this in agreement with (Barkosky and Einhellig, 1993).

Conclusion

From this study we can be concluded that priming of faba bean with low concentrations of SA (0.5 mM) is a simple, cheap and unsophisticated tool that has a practical importance and could be recommended to farmers to achieve higher germination and uniform emergence under field conditions to accelerate potential crop yield in near future. In future, the exogenous application of this phytohormone might act as powerful tool in enhancing the growth and productivity of more biomass and photosynthetic capacity, especially in areas with saline irrigation water resources at planting stage. Moreover; the planted seeds will be less susceptible to soil-borne pests and diseases, and this has positive consequences on management of this crop at planting and germination stage.

Acknowledgments

Authors are thankful to Heads of Biology and Chemistry Departments; also with deep grateful thanks to Mrs. Mariam; Mrs. Sarah and Mrs. Asmaa laboratory technicians in Biology and Chemistry Departments, Faculty of Science, Yanbu, Taibah University, Saudi Arabia for providing necessary facilities to carry out the present work.

References

- Abd El-Wahed, M. S. A., Amin, A. A. and El-Sh, M. R. (2006). Physiological effect of some bioregulators on vegetative growth, yield and chemical constituents of yellow maize plants. *World Journal of Agricultural Sciences* 2:149-155.
- Abdul-Baki, A. A. and Anderson, J. D. (1973). Vigour determination of Soybean seed by multiple criteria. *Crop Science* 13:630-633.
- Ahmad, J., Khaliq, T., Ahmad, A., Basra, S., Hasnain, Z. and Ali, A. (2012). Effect of seed priming with ascorbic acid, salicylic acid and hydrogen peroxide on emergence, vigor and antioxidant activities of maize. *African Journal of Biotechnology* 11:1127-1132.
- Ansari, O. and Sharif-Zadeh, F. (2012). Osmo and hydro priming improvement germination characteristics and enzyme activity of Mountain Rye (*Secale montanum*) seeds under drought stress. *Journal of Stress Physiology and Biochemistry* 8:253-261.
- Barkosky, R. R. and Einhelling, F. A. (1993). Effect of salicylic acid on plant water relationship. *Journal of Chemical Ecology* 19:237-247.
- Bezaukova, M., Sakhabutdinova, V. and Fatkhutdinova, F. A. R. (2001). The role of hormonal changes in protective action of salicylic acid on growth of wheat seedlings under water deficit. *Agrochemicals* 2:51-54.
- Blokhina, O., Virolainen, E. and Fagerstedt, K. V. (2003). Antioxidants, oxidative damage and oxygen deprivations stress. *Annals of Botany* 91:179- 194.
- Delavari, P. M., Baghizadeh, A., Enteshari, S. H., Kalantari, K. M., Yazdanpanah, A. and Mousavi, E. A. (2010). The Effects of salicylic acid on some of biochemical and morphological characteristic of *Ocimum basilicum* under salinity stress. *Australian Journal of Basic and Applied Sciences* 4:4832-4845.
- Durner, J. and Klessig, D. F. (1995). Inhibition of ascorbate peroxidase by salicylic acid and 2,6-dichloroisonicotinic acid, two inducers of plant defense responses. *Proceedings of the National Academy of Sciences* 92:11312-11316.
- Eastmond, P. J. and Graham, I. A., (2001). Re-examining the role of the glyoxylate cycle in oilseeds. *Trends in Plant Science* 6:72-8.
- Eisvand, H. R., Shahrosvand, S., Zahedi, B., Heidari, S. and Afroughe, S. (2011). Effects of hydro-priming and hormonal priming by gibberellin and salicylic acid on seed and seedling quality of carrot (*Daucus carota* var. *sativus*). *Iranian Journal of Plant Physiology* 1:233-239.
- El-Khallal, S. M., Hathout, T. A., Abd El Raheim, A., Ashour, A. and Kerrit, A. A. (2009). Brassinolide and salicylic acid induced growth, biochemical activities and productivity of maize plants grown under salt stress. *Research journal of agriculture and biological sciences* 5:380-390.
- El-Tayeb, M. A. (2005). Response of barley grains to the interactive effects of salinity and salicylic acid. *Plant Growth Regulation* 45:215-224.

- Farahbakhsh, H. (2012). Germination and seedling growth in un-primed and primed seeds of Fenel as affected by reduced water potential induced by NaCl. *International Research Journal of Applied and Basic Sciences* 3:737-744.
- Gunes, Y., Inal, A., Alpaslan, M., Eraslan, F., Bagci, E. G. and Cicek, G. N. (2007). Salicylic acid induced changes on some physiological parameters symptomatic for oxidative stress and mineral nutrition in maize (*Zea mays* L.) grown under salinity. *Journal of Plant Physiology* 164:728-736.
- Habibi, A. and Abdoli, M. (2013). Influence of salicylic acid pre-treatment on germination, vigor and growth parameters of garden cress (*Lepidium sativum*) seedlings under water potential loss at salinity stress. *International Research Journal of Applied and Basic Sciences* 4:1393-1399.
- Hermann, K., Meinhard, J., Dobrev, P., Linkies, A., Pesek, B., Heß, B. and Leubner-Metzger, G. (2007). 1-Aminocyclopropane-1-carboxylic acid and abscisic acid during the germination of sugar beet (*Beta vulgaris* L.): a comparative study of fruits and seeds. *Journal of Experimental Botany* 58:3047-3060.
- Hopkins, W. G. and Huner, N. P. A. (2004). *Introduction to plant physiology*. 3rd ed.
- ISTA (1985). *International rules for seed testing*. *Seed Science and Technology* 13:361-513.
- Jam, B. J., Shekari, F. A. R. I. D., Azimi, M. R. and Zangani, E. S. M. A. E. I. L. (2012). Effect of priming by salicylic acid on germination and seedling growth of safflower seeds under CaCl₂ stress. *International Journal of Agricultural Research* 2:1097-1105.
- Klessig, D. F., Vlot, C. A. and Dempsey, D. A. (2009). Salicylic acid, a multifaceted hormone to combat disease. *Annual Review Phytopathology* 47:177-206.
- Ma, G., Jin, Y., Piao, J., Kok, F., Bonnema, G. and Jacobsen, E. (2005). Phytate, calcium, iron and zinc contents and their molar ratios in food commonly consumed in China. *Journal of Agricultural and Food Chemistry* 53:10285-10290.
- Maia, F. C., Moraes, D. M. and Moraes, R. C. P. (2000). Atividade total da fosfatase ácida e da α -amilase induzidas por ácido jasmônico em sementes de soja. *Revista Brasileira de Sementes* 22:259-263.
- Mazaheri-e-Tirani, M. and Manouchehri, K. (2005). Effects of the role of salicylic acid, drought stress, ethylene and interaction of three factors on seed germination of *Brassica napus*. *Iranian Journal of Biology* 19:408-418.
- Metraux, J. P., Signer, H., Ryals, J., Ward, E., Wyss-Benz, M., Gaudin, J., Raschdorf, K., Schmid, E., Blum, W. and Inverardi, B. (1990). Increase in salicylic acid at the onset of systemic acquired resistance in cucumber. *Science* 250:1004-1006.
- Metwally, A., Finkemeier, I., Georgi, M. and Dietz, K. J. (2003). Salicylic acid alleviates the cadmium toxicity in barley seedlings. *Plant Physiology* 132:272-281.
- Nun, N.B., Plakhine, D., Joel, D. M. and Mayer, A. M. (2003). Changes in the activity of the alternative oxidase in Oribanche seeds during conditioning and their possible physiological function. *Photochemistry* 64:235-241.

- Rajjou, L., Belghazi, M., Huguet, R., Robin, C., Moreau, A., Job, C. and Job, D. (2006). Proteomic investigation of the effect of salicylic acid on Arabidopsis seed germination and Establishment of early defense mechanism. *Plant Physiology* 141:910-923.
- Rao, M. V., Paliyath, G., Ormrod, D. P., Murr, D. P. and Watkins, C. B. (1997). Influence of salicylic acid on H₂O₂ production, oxidative stress, and H₂O₂-metabolizing enzymes. *Plant Physiology* 115:137-149.
- Raskin, I. (1992). Role of salicylic acid in plants. *Plant Molecular Biology* 43:439-463.
- Russell, J. W., Golovoy, D., Vincent, A. M., Mahendru, P., Olzmann, J. A. and Mentzer, A. (2002). High glucose induced oxidative stress and mitochondrial dysfunction in neurons. *The FASEB Journal* 16:1738-1748.
- Shah, J. (2003). The salicylic acid loop in plant defense. *Current opinion in plant biology* 6:365-371.
- Shakirova, F. M., Sakhabutdinova, A. R., Bezrukova, M. V., Fathudinova, R. A. and Fathudinova, D. R. (2003). Changes in hormonal status of wheat seedlings induced by Salicylic acid and salinity. *Plant Science* 164:317-322.
- Sharafizad, M., Naderi, A., Siadat, S. A., Sakinejad, T. and Lak, S. (2013). Effect of Salicylic Acid Pretreatment on Germination of Wheat under Drought Stress. *Journal of Agricultural Science* 5:179-199.
- Shehata, S. A. M., Ibrahim, S. I. and Zaghlool, S. A. M. (2001). Physiological response of flag leaf and ears of maize plant induced by foliar application of kinetin, KIN and acetyl salicylic acid, ASA. *Annals of Agricultural Science, Ain Shams University* 46:435-449.
- Shi, Q. and Zhu, Z. (2008). Effects of exogenous salicylic acid on manganese toxicity, element contents and antioxidative system in cucumber. *Environmental and Experimental Botany* 63:317-326.
- Singh, B. and Usha, K. (2003). Salicylic acid induced physiological and biochemical changes in wheat seedlings under water stress. *Plant Growth Regulation* 39:137-141.
- Singh, K., Chaturvedi, P., Kumar, V. and Bose, B., (2010). Effects of salicylic acid on seedling growth and nitrogen metabolism in cucumber (*Cucumis sativus* L.). *Journal of Stress Physiology and Biochemistry* 6:103-113.
- Szepesi, Á., Csiszár, J., Bajkán, Sz., Gémes, K., Horváth, F., Erdei, L., Deér, A., Simon, L. M. and Tari, I. (2005). Role of salicylic acid pre-treatment on the acclimation of tomato plants to salt and osmotic stress. *Acta Biologica Szegediensis* 49:123-125.
- Van Camp, W., Van Montagu, M. and Inzé, D. (1998). H₂O₂ and NO: redox signals in disease resistance. *Trends in Plant Science* 3:330-334.
- Wu, L., Guo, X. and Harivandi, M.A., (1998). Allelopathic effects of phenolic acids detected in buffalograss (*Buchloe dactyloides*) clippings on growth of annual bluegrass (*Poa annua*) and buffalograss seedlings. *Environmental and Experimental Botany* 39:159-167.

- Yalpani, N., Enyedi, A. J., León, J. and Raskin, I. (1994). Ultraviolet light and ozone stimulate accumulation of salicylic acid, pathogenesis-related proteins and virus resistance in tobacco. *Planta* 193:372-376.
- Zaghlool, S. A. M. (2002). The effect of gibberellic acid (GA3), salicylic acid (SA), spermidine (Spd) and methods of application on growth, yield, some chemical constituents and some phytohormones in mungbean (*Vigna radiata* L.). *Arab Universities Journal of Agricultural Sciences* 2:493-504.

(Received: 10 October 2016, accepted: 5 November 2016)