
Study for Probiotic Properties of *Lactobacillus salivarius* KL-D4 Isolated from Duck Intestine

Rumjuankiat, K.¹, Ngamyeesoon, N.², Swetwiwathana, A.³ and Pilasombut, K.^{4*}

¹Faculty of Biotechnology, College of Agriculture Innovation Biotechnology and Food, Rangsit University, LakHok, MueangPathumThani District, PathumThani 12000, Thailand; ²Division of Plant Production Technology, Faculty of Agricultural Technology, King Mongkut's Institute of Technology Ladkrabang, Bangkok, 10520, Thailand.; ³Division of Fermentation Technology, Faculty of Agro-Industry, King Mongkut's Institute of Technology Ladkrabang, Bangkok 10520, Thailand; ⁴Department of Animal Production Technology and Fisheries, Faculty of Agricultural Technology, King Mongkut's Institute of Technology Ladkrabang, Bangkok, 10520, Thailand.

Rumjuankiat, K., Ngamyeesoon, N., Swetwiwathana, A. and Pilasombut, K. (2017). Study for probiotic properties of *Lactobacillus salivarius* KL-D4 isolated from duck intestine. International Journal of Agricultural Technology 13(6):823-837.

Abstract *Lactobacillus salivarius* KL-D4 isolated from duck intestine, could produce bacteriocin against several pathogenic and spoilage bacteria. The study aimed to evaluate the probiotic potential of *L.salivarius* KL-D4 as preliminary study for probiotic properties. The survival of *L.salivarius* KL-D4 was performed in 1-5%NaCl concentration, pH range of 2-10 and bile salt concentrations at 0.3%, 0.6% and 0.9%. The survival rate of this strain was determined at pH 2, 3, 4 and 7 in simulated gastrointestinal tract. Furthermore, the susceptibility to antibiotic was also investigated. It was found that this strain could tolerate to NaCl (1-5%), pH 3-10 and bile salt (0.3-0.9%). Additionally, it was able to survive in simulated gastric juice and showed the survival rate approximately 42%, 85% and 98% at pH 3, 4 and 7, respectively. The tolerance of *L. salivarius* KL-D4 in intestinal fluids at pH 3, 4 and 7 with a viability count about log 2.0, 4.8 and 6.8 CFU/ml, respectively. Moreover, this strain was also sensitive to ampicillins, chloramphenicol, cephalothin, erythomycin, nitrofurantoin, sulfamethoxazole/trimethoprim, penicillin G and amoxicillin. Therefore, *L. salivarius* KL-D4 can be probiotic alternative in the future.

Keywords: *Lactobacillus salivarius*, Probiotic, Duck Intestine

Introduction

Lactic acid bacteria (LAB) are defined as a cluster of lactic acid producing, non spore forming, low G+C content and Gram-positive bacteria (Gómez *et al.*, 2016). They can produce antibacterial substance and are considered as generally recognized as safe (GRAS), resulting in a great

* **Coressponding author:** Pilasombut, K.; **Email:** komkhae.pi@kmitl.ac.th.

biotechnological potential in the food industry (Alvarez-Sieiro *et al.*, 2016). LAB is widely used as starter culture including cheeses, yoghurts, fermented milk (Hoque *et al.*, 2010), Chinese fermented cabbage (Zhao *et al.*, 2016), kimchi (Jung *et al.*, 2012), fermented oyster mushrooms (Liu *et al.*, 2016) and Thai traditional fermented sausage such as Nham (Kingchaet *et al.*, 2012, Swetwivathana and Visessanguan, 2015), Plaa-som (Saithong *et al.*, 2010) and Som-fug (Kongkiattikajorn, 2015). Furthermore, LAB can be used as biopreservative for creamy filling (Therdathaet *et al.*, 2016) or used as sanitizer (Rumjuankiat *et al.*, 2017). In addition, LAB are commonly studied as probiotic for the past few decades because they are desirable microflora of the gastrointestinal tract (Angmo *et al.*, 2016).

Probiotics are living microorganisms that provide health benefits to host (FAO/WHO, 2014). Generally, probiotics should possess some properties such as contributing colonization resistance, supporting the intestinal barrier including tight junction expression or secretion of mucus, instructing the intestinal microbiota organization and activity (Gómez *et al.*, 2016). Thus, the functional requirements of probiotics is necessary to investigate such as tolerance to acid, resistance to gastric acidity and bile salts or production of antimicrobial compound that comprised lactic acid, acetic acid, hydrogen peroxide or bacteriocins (Shehata *et al.*, 2016).

The genus *Lactobacillus* belongs to the large group of LAB and widely used in food both as starters in fermented food and probiotics (Arena *et al.*, 2016). Among lactobacilli, *Lactobacillus salivarius* have been extensively used for probiotic. It is a promising probiotic usually isolated from human feces (Martín *et al.*, 2006, Tinrat *et al.*, 2011), human milk (Langaet *et al.*, 2012), mammalian digestive tract (Neville and O'Toole, 2010), or avian gastrointestinal tracts (Kergourlay *et al.*, 2012, Nouri *et al.*, 2010). Many of *Lactobacillus salivarius* are producers of unmodified bacteriocins of subclasses IIa, IIb and IIc (O'Shea *et al.*, 2011). Moreover, there are a lot of mechanisms of *Lactobacillus salivarius* enhance intestinal health (Messaudi *et al.*, 2013).

In previous study, Therdathaet *et al.* (2016) have been report about *Lactobacillus salivarius* KL-D4, which was a new strain isolated from duck intestine, could produce bacteriocin called salivaricin KLD. Its partial purified bacteriocin was tested on the artificial contamination of creamy filling and showed the inhibition against *Bacillus cereus*, *Enterococcus faecalis*, *Pseudomonas stutzeri*, *Staphylococcus* sp. and *Stenotrophomonas* sp. It seems that, salivaricin KLD might be a possible biopreservative for food industry. Therefore, the present study aimed on evaluation of potential bacteriocin-

producing strains, *Lactobacillus salivarius* KL-D4, to investigate various probiotic properties.

Materials and methods

Bacterial strains and growth conditions

Lactobacillus salivarius KL-D4 was cultivated in MRS medium (de Man, Rogosa and Sharpe, Merck, Germany) at 30 °C for 16 h without agitation. Indicator strains used for the antibacterial assay were propagated according to the growth conditions shown in Table 1.

Table 1. List of indicator strains and their growth conditions

| Indicator strains | Medium | Temperature (°C) |
|---|---------|------------------|
| Lactic acid bacteria | | |
| <i>Lactobacillus sakei</i> subsp. <i>sakei</i> JCM 1157 ^T | MRS | 30 |
| <i>Pediococcus dextrinicus</i> JCM 5887 ^T | MRS | 30 |
| <i>Lactobacillus plantarum</i> ATCC 14917 | MRS | 30 |
| <i>Enterococcus faecium</i> JCM 5804 ^T | MRS | 30 |
| Other Gram-positive bacteria | | |
| <i>Listeria innocua</i> ATCC 33090 ^T | TSB-YE | 37 |
| <i>Bacillus coagulans</i> JCM 2257 ^T | TSB-YE* | 37 |
| <i>Bacillus subtilis</i> subsp. <i>subtilis</i> JCM 1465 ^T | TSB-YE* | 37 |
| Gram-negative bacteria | | |
| <i>Salmonella enterica</i> serovar <i>Enteritidis</i> DMST 17368 | TSB-YE* | 37 |
| <i>Salmonella</i> Typhimurium TISTR 292 | TSB-YE* | 37 |
| <i>Pseudomonas fluorescens</i> TISTR 358 | TSB-YE* | 26 |
| <i>Pseudomonas fluorescens</i> JCM 5963 ^T | TSB-YE* | 26 |
| <i>Escherichia coli</i> O157:H7 | NB* | 37 |

ATCC, American Type Culture Collection, Rockville, Md, USA; JCM, Japan Collection of Microorganisms, Wako, Japan; TISTR, Thailand Institute of Scientific and Technological Research; MRS, De Man, Rogosa and Sharpe broth (Merck, Germany); TSB-YE, Tryptic Soy Broth (Merck, Germany) containing 0.6% Yeast extract (Merck, Germany); NB, Nutrient broth (Merck, Germany)

* Growth under agitation at 200 rpm.

Antibacterial activity assay

Bacterial inhibition

The pathogenic and spoilage bacteria inhibition by *L. salivarius* KL-D4 was determined using agar well diffusion assay (Ranganath and Sharmila, 2016). The pathogenic and spoilage bacteria including *Salmonella*

*entericaserovar*Enteritidis DMST 17368, *Salmonella*Typhimurium TISTR 292 and *Escherichia coli* O157:H7 were used as indicators (Table 1). Each indicator strain was lawn MRS agar plate using sterile cotton swab and incubated for 15 min. The culture of *L. salivarius* KL-D4 was centrifuged at 12,000 rpm for 10 min. Subsequently, 50 µl of the cell free supernatant (CFS) was transferred into agar well which was made using sterile cork borer and incubated overnight. The diameter of the inhibition zone was measured.

Antimicrobial inhibition

The antimicrobial substances of *L. salivarius* KL-D4 was determined for its inhibition activities against 9 indicators which were listed in Table 1. MRS agar plate was overlaid with 7 ml of MRS soft agar (1% agar, w/v) mixed with 30 ml of each indicator (approximately 10^6 CFU/ml). After setting, 10 µl of the CFS was spotted onto the surface and incubated overnight at the optimum temperature. The inhibition zone was determined using the spot-on-lawn method according to the modified method of Ennahar *et al.* (2001).

Investigation of L. salivarius KL-D4 tolerance

pH tolerance

This method was modified according to the method of Hoque *et al.* (2010). Two percent of fresh *L. salivarius* KL-D4 was transferred to 10 ml of MRS broth adjusted to different pH at 2.0, 3.0, 4.0, 5.0, 6.0, 7.0, 8.0, 9.0 and 10.0 with 0.5 mol l^{-1} HCl or 0.5 mol l^{-1} NaOH, and incubated at 37°C for 18 hr. After incubation, the growth was investigated by observing the turbidity. Double positive sign (++), positive sign (+) and negative sign (-) was indicated for maximum growth, normal growth and no growth, respectively.

NaCl tolerance

The NaCl tolerance was investigated by preparing MRS broth containing various concentrations (0%, 1%, 2%, 3%, 4%, 5%, 6% and 7%) of NaCl. Two percent of fresh *L. salivarius* KL-D4 was inoculated in each test tube and incubated at 37°C for 18 hr. MRS broth without NaCl was used as a control. All samples were incubated at 37°C for 18 hr. The turbidity of the maximum growth were determined as double positive sign (++), normal growth as a positive sign (+) while no growth as a negative sign (-) according to the modified method of Hoque *et al.* (2010).

Bile tolerance

The culture was grown in MRS broth at 30°C for 18 hr then inoculated into 10 ml of MRS broth contained bile salts (Sigma-Aldrich, St. Louis, MO, USA) at various concentration (0%, 0.3%, 0.6% and 0.9%). All samples were incubated at 30°C for 18 hr. Subsequently, 100 µl of each sample was spread onto MRS agar and incubated at 30°C for 18 hr under anaerobic condition. Tolerance to bile salts were assayed by growing culture in MRS plate in duplicate. The method used for testing was similar to the modified method of Bakari *et al.* (2011).

In vitro simulated gastric juice and bile salt solution

The capability of *L. salivarius* KL-D4 to tolerate the simulated gastric juice and bile salt solution was determined as described by Zárate *et al.* (2000). Two percent of *L. salivarius* KL-D4 was inoculated in 100 ml of MRS broth and incubated at 30°C for 16 hr. The culture was centrifuged at 4,000 rpm, 4°C for 10 min and washed in sterile saline buffer (NaCl, 0.9% w/v). Then, dissolved the cell in 100 ml of artificial gastric juice (125mM NaCl, 7mM KCl, 45 mM NaHCO₃, 3% pepsin adjusted the final pH to 2, 3, and 4 with HCl). The cell survival was considered in simulated gastric juice by standard plate count. To simulate peristalsis and the normal temperature of human body, the suspension was incubated at 37°C with agitation (200 rpm.) for 30, 60, 90 and 180 min. An aliquot of the suspension was enumerated for survival cell. Subsequently, 50 ml of the suspension was centrifuged at 4,000 rpm, 4°C for 10 min, discarded the supernatant and suspended the pellets in 50 ml of intestinal fluid including 0.1% pancreatin USP (Sigma-Aldrich, St. Louis, MO, USA) and 0.15% bile salts (Sigma-Aldrich, St. Louis, MO, USA) adjusted pH to 8.0 with 5 N NaOH. The suspensions were incubated for 0, 30, 60, and 180 min. The cell survival was calculated equation below:

$$\% \text{bacterial survival} = N_{\text{survival}} \times 100 / N_{\text{initial}}$$

Where N_{survival} is the number of bacterial survival at 0, 30, 60, 90 and 180 min (log cfu/ml) and N_{initial} is the number of initial cell (log cfu/ml).

Safety assessment

The antibiotic resistance was investigated using agar overlay disc diffusion test according to the modified method of Tulumoglu *et al.* 2013. *L. salivarius* KL-D4 was propagated in 10 mL MRS broth at 30 °C overnight.

After incubation, the culture was adjusted to 0.5 McFarland and swabbed onto MRS agar plate. Subsequently, antibiotic discs (Oxoid, England) such as Ampicillins (10 µg), Chloramphenicol (30 µg), Cephalothin (30 µg), Erythromycin (15 µg), Gentamycin(10 µg), Kanamycin (30 µg), Naldixic acid (30 µg), Neomycin (30 µg), Nitrofurantoin (300 µg), Norfloxacin (10 µg), Novabincin (5 µg), Oxolinicacid (2 µg), Tetracyclin (30 µg), Sulfamethoxazole/Trimethoprim (25 µg/1.5 µg)andOxytetracyclin (30 µg) were placed on the swabbed plates with sterile conditions. After 24 h incubation at 30 °C, the diameter of inhibition zone was measured for its susceptibility. The interpretation were designated by the CLSI (Ferraro, 2001).

Results

Inhibition spectra of CFS produced by L. salivarius KL-D4

The antimicrobial substance of *L. salivarius* KL-D4 were determined against three pathogenic bacteria, five food spoilage and related four LABs shown in Table 2. The CFS of *L. salivarius* KL-D4 against both potential Gram-positive and Gram-negative bacteria. Especially, *Salmonella enterica* serovar Enteritidis DMST 17368, *Salmonella* Typhimurium TISTR 292 and *Escherichia coli* O157:H7, which were Gram-negative bacteria pathogens, were shown with inhibition zone approximately 15, 19 and 16 mm, respectively. The most sensitive strains to KL-D4 was *L. sakei* subsp. *sakei* JCM 1157^T, whereas *P. dextrinicus* JCM 5887^T was the most resistant.

Table 2. Antimicrobial spectrum of *L. salivarius* KL-D4 against pathogenic bacteria, food spoilage and related LAB

| Indicator strains | Inhibition zone (mm) |
|---|----------------------|
| <i>Lactobacillus sakei</i> subsp. <i>sakei</i> JCM 1157 ^T | 25 |
| <i>Pediococcus dextrinicus</i> JCM 5887 ^T | 6 |
| <i>Lactobacillus plantarum</i> ATCC 14917 | 13 |
| <i>Enterococcus faecium</i> JCM 5804 ^T | 12 |
| <i>Listeria innocua</i> ATCC 33090 ^T | 22 |
| <i>Bacillus coagulans</i> JCM 2257 ^T | 12 |
| <i>Bacillus subtilis</i> subsp. <i>subtilis</i> JCM 1465 ^T | 20 |
| <i>Salmonella enterica</i> serovar Enteritidis DMST 17368 | 15 |
| <i>Salmonella</i> Typhimurium TISTR 292 | 19 |
| <i>Pseudomonas fluorescens</i> TISTR 358 | 16 |
| <i>Pseudomonas fluorescens</i> JCM 5963 ^T | 17 |
| <i>Escherichia coli</i> O157:H7 | 16 |

pH, NaCl and bile tolerance

The effect of various pH, NaCl and bile salt concentrations on the viability of *L. salivarius* KL-D4 was shown in Table 3. This strain could survive in wide range conditions of pH (3-10), NaCl (1%-5%) and bile salt (0.3%-0.9%). Moreover, *L. salivarius* KL-D4 well growth at pH 6-8, 1%-2% of NaCl and 0.3% bile salt concentrations compared with control sample.

Table 3. The viability of *L. salivarius* KL-D4 under various pH, NaCl and bile salt concentrations.

| Treatments | Growth level |
|-------------------------|--------------|
| pH value | |
| 2.0 | - |
| 3.0 | + |
| 4.0 | + |
| 5.0 | + |
| 6.0 | ++ |
| 7.0 | ++ |
| 8.0 | ++ |
| 9.0 | + |
| 10.0 | + |
| NaCl concentration | |
| 0% | ++ |
| 1% | ++ |
| 2% | ++ |
| 3% | + |
| 4% | + |
| 5% | + |
| 6% | - |
| 7% | - |
| Bile salt concentration | |
| 0% | ++ |
| 0.3% | ++ |
| 0.6% | + |
| 0.9% | + |

Legend: ++, maximum growth; +, normal growth; -, no growth

Survival of bacteria in simulated gastric digestion

The survival of *L. salivarius* KL-D4 in artificial gastric and intestinal fluid was evaluated. This strain could not survive in simulated gastric at pH 2 after inoculation, on the contrary it could survive for 180 min at pH 3, 4 and 7, with a viability count about log 3.7, 6.7 and 7.8 CFU/ml, respectively, as shown

in Figure 1A. Furthermore, it showed a survival rate approximately 42%, 85% and 98% at pH 3, 4 and 7, respectively. In addition, *L. salivarius* KL-D4 tolerated to intestinal fluids at pH 3, 4 and 7 with a viability count about log 2.0, 4.8 and 6.8 CFU/ml, respectively. After 180 min, the viability count of *L. salivarius* KL-D4 in intestinal fluid decreased from a various viability count in gastric digestion to log 2, 4.8 and 6.8 CFU/ml and the survival rate indicated about 25%, 60.8% and 84.9% at pH 3, 4 and 7, respectively (Figure 1B).




Figure 1. The viable count (log cfu/ml) and survival percentage (%) of *L. salivarius* KL-D4 determined by (A) simulated gastric and (B) intestinal fluid at 0, 30, 60, 90 and 180 min. The viable counts was evaluated at pH 2 (—◆—), pH 3 (—■—), pH 4 (—●—) and pH 7 (—▲—). The survival percentage was investigated at pH 2 (···◇···), pH 3 (···□···), pH 4 (···○···) and pH 7 (···△···).

Antibiotic resistance

Antibiotic susceptibility of *L. salivarius*KL-D4 was investigated according to the anti-microbial drug sensitivity standard of CLSI criteria. There were 18 antibiotics for the sensitivity test as shown in Table 4. *L. salivarius*KL-D4 was generally susceptible to 8 antibiotics including ampicillins, chloramphenicol, cephalothin, erythomycin, nitrofurantoin, sulfamethoxazole/trimethoprim, penicillin G and amoxycillin. On the other hand, *L. salivarius*KL-D4 could resist to tetracyclin, especially gentamycin, kanamycin, naldixic acid, neomycin, norfloxacin, oxolinic acid, oxytetracyclin and streptomycin were extremely resisted by this strain.

Table 4, Antimicrobial susceptibility of *L. salivarius* KL-D4

| Antimicrobial agents | Disk content (µg) | Zone diameter (mm) | Acceptable inhibitory |
|-------------------------------|--------------------------|---------------------------|------------------------------|
| Ampicillins | 10 | 30 | S |
| Chloramphenicol | 30 | 29 | S |
| Cephalothin | 30 | 30 | S |
| Erythomycin | 15 | 25 | S |
| Gentamycin | 10 | 0 | R |
| Kanamycin | 30 | 0 | R |
| Naldixic acid | 30 | 0 | R |
| Neomycin | 30 | 0 | R |
| Nitrofurantoin | 300 | 19 | S |
| Norfloxacin | 10 | 0 | R |
| Novabincin | 5 | 15 | I |
| Oxolinic acid | 2 | 0 | R |
| Tetracyclin | 30 | 9 | R |
| Sulfamethoxazole/Trimethoprim | 25/1.5 | 20 | S |
| Oxytetracyclin | 30 | 0 | R |
| Penicillin G | 10 | 30 | S |
| Amoxycillin | 10 | 30 | S |
| Streptomycin | 10 | 0 | R |

Discussion

Although many researchers have been reported about the probiotic properties and safety using lactobacilli, it should not ignore the assessment of appropriate *in vitro* assays (Borriello *et al.*, 2003, Messaoudi *et al.*, 2013, Sanders *et al.*, 2010), especially *Lactobacillus sarivarius*. Several *L.sarivarius* strains comprised *L.sarivarius*(Martín *et al.*, 2006), *L.sarivarius*W24(Koninkx *et al.*, 2010), *L. salivarius* C3 (Kirtzalidou *et al.*, 2011) and *L.*

salivarius REN(Wang *et al.*, 2017) were considered as potential probiotics. Not only *L. salivarius* was proved as probiotics, but some strain of *L. salivarius* also can produce bacteriocin such as *L. salivarius* K7 (Narakaew *et al.*, 2010, Pilasombut *et al.*, 2006), *L. salivarius* subsp. *salivarius* UCC118 (Flynn *et al.*, 2007), *L. salivarius* DPC6005 (Walsh *et al.*, 2008) or *L. salivarius* SMXD51 (Saint-Cyr *et al.*, 2017). In previous study, *L. salivarius* KL-D4 was confirmed that it could produce bacteriocin and consequently showed the activity of microbial inhibition (Therdthatha *et al.*, 2016). However, bacteriocin of *L. salivarius* KL-D4, which was neutralized, could not inhibit *E. coli* O157:H7 while its CFS without pH adjustment showed the inhibition of this pathogenic strain in this experiment. Therefore, our results demonstrated that *L. salivarius* KL-D4 may have good probiotic because it against both Gram positive and Gram negative bacteria. Moreover, *L. salivarius* KL-D4 could inhibit all tested pathogenic bacteria, especially *Escherichia coli*, *Salmonella* Typhimurium or *Staphylococcus aureus*. These pathogenic bacteria were occasionally found in gastrointestinal tract and might cause gastroenteritis (Klayraung *et al.*, 2008).

NaCl is an inhibitory substance because it could affect physiology, enzymatic, water activity and metabolism of the bacterial cell when it was cultivated in a high salt concentration (Ibourahema *et al.*, 2008, Liu *et al.*, 1998, Menconi *et al.*, 2014). LAB strains with high osmotolerance would be a requirement strain (Ibourahema *et al.*, 2008) because when LAB produced lactic acid, alkali would be pumped into the broth to prevent an excessive reduction in pH. Therefore, the free acid would be converted to its salt form and then increasing the osmotic pressure on the bacterial cells (Menconi *et al.*, 2014). In our results, *L. salivarius* KL-D4 was able to tolerate NaCl concentration approximately 1-5% and supposed that could be a high osmotolerance strain, thereby resulting in the strain would be desirable as a commercial strain.

pH is a crucial factors that affect bacterial growth. However, some LAB strains can grow in both acidic and alkaline (Hoque *et al.*, 2010). In our experiment, the effect of pH ranging from 2 to 10 on the survival strain of *L. salivarius* KL-D4 was studied. This strain could survive in pH 3-5 and the growth could dramatically rise from pH 6 to pH 8. Subsequently, the growth decreased when it was cultured in pH 9-10. However, the potential probiotics should resist to acid at least pH 3.0 (Fernández *et al.*, 2003, Sahadeva *et al.*, 2011) because bacterial strain have to tolerate for gastric stress and could survive for longer period in food which have a high acid content without larger reduction in humans (Shehata *et al.*, 2016).

Not only the ability of survival cell in low pH condition, but the capability to tolerate the high concentration of bile salt of the upper gastrointestinal tract

is considered as good indicators also (Kizerwetter-Świda and Binek, 2016). Normally, the maximum concentration of bile salt in intestinal tract of healthy men is around 0.3% (Hoque *et al.*, 2010) while the concentration of bile salt in the small intestine is approximately 0.2%-2.0% (Klayraung *et al.*, 2008). In this experiment, the concentrations of 0.3%, 0.6% and 0.9% bile salts were used. Our strain, *L. salivarius* KL-D4, was able to tolerate up to 0.9% of bile salts concentration.

However, the ability of bacterial survival when they pass through the stomach is the first requirement for probiotics (Fernández *et al.*, 2003). The viability and survival rates at pH 2 of *L. salivarius* KL-D4 in gastric is generally low (less than 1%) and could not survive in intestinal fluid. The viability and survival rate of *L. salivarius* KL-D4 was gradually decreased from 7.72 to 3.36 log CFU/ml and 93.37% to 42.05% at pH 3 for 180 min, respectively. The viable cell count in intestinal fluid was about 2 log CFU/ml and showed calculated survival rate about 25% at 180 min. These results indicated that there were the viability of cell in simulated gastrointestinal condition. Although *L. salivarius* KL-D4 demonstrated that it could not tolerate to pH 2 in gastric and intestinal fluid, it might be survive when consumed in food or encapsulated using different biopolymeric substances (Ashraf and Smith, 2016, Chávarri *et al.*, 2010).

In our experiment, *L. salivarius* KL-D4 was susceptible to ampicillins, chloramphenicol, cephalothin, erythromycin, nitrofurantoin, sulfamethoxazole/trimethoprim, penicillin G and amoxicillin. Another criterion to be a good source of probiotics is antibiotic susceptibility. This is a crucial point because bacteria, which used as probiotics, may serve as host of antibiotic resistance gene, resulting in transferred to pathogenic bacteria (Klayraung *et al.*, 2008).

In conclusion, *L. salivarius* KL-D4 exhibits broad spectrum inhibition and tolerate to 1-5% NaCl, pH 3-10, 0.3%-0.9% bile salt concentration and also survive in simulated gastric digestion. In addition, this strain sensitive to ampicillins, chloramphenicol, cephalothin, erythromycin, nitrofurantoin, sulfamethoxazole/trimethoprim, penicillin G and amoxycillin. Therefore, *L. salivarius* KL-D4 might be a potential probiotic strains in the future.

Acknowledgement

This research is partially supported by Meat Technology Research Network Center of King Mongkut's Institute of Technology Ladkrabang on corresponding between King Mongkut's Institute of Technology Ladkrabang and the Thailand Research Fund.

References

- Alvarez-Sieiro, P., Montalbán-López, M., Mu, D. and Kuipers, O. P. (2016). Bacteriocins of lactic acid bacteria: extending the family. *Applied Microbiology and Biotechnology* 100:2939-2951.
- Angmo, K., Kumari, A., Savitri, T. and Bhalla, T. C. (2016). Probiotic characterization of lactic acid bacteria isolated from fermented foods and beverage of Ladakh. *LWT - Food Science and Technology* 66:428-435.
- Arena, M. P., Silvain, A., Normanno, G., Grieco, F., Drider, D., Spano, G. and Fiocco, D. (2016). Use of *Lactobacillus plantarum* strains as a bio-control strategy against food-borne pathogenic microorganisms. *Frontiers in Microbiology* 7:464.
- Ashraf, R. and Smith, S. C. (2016). Commercial lactic acid bacteria and probiotic strains-tolerance to bile, pepsin and antibiotics. *International Food Research Journal* 23:777-789.
- Bakari, D., Tatsadjieu, N. L., Mbawala, A. and Mbofung, C. M. (2011). Assessment of physiological properties of some lactic acid bacteria isolated from the intestine of chickens use as probiotics and antimicrobial agents against enteropathogenic bacteria. *Innovative Romanian Food Biotechnology* 8:33-40.
- Borriello, S. P., Hammes, W. P., Holzapfel, W., Marteau, P., Schrezenmeir, J., Vaara, M. and Valtonen, V. (2003). Safety of probiotics that contain lactobacilli or bifidobacteria. *Clinical Infectious Diseases* 36:775-780.
- Chávarri, M., Marañón, I., Ares, R., Ibáñez, F. C., Marzo, F. and Villarán, M.d. C. (2010). Microencapsulation of a probiotic and prebiotic in alginate-chitosan capsules improves survival in simulated gastro-intestinal conditions. *International Journal of Food Microbiology* 142:185-189.
- Ennahar, S., Asou, Y., Zendo, T., Sonomoto, K. and Ishizaki, A. (2001). Biochemical and genetic evidence for production of enterocins A and B by *Enterococcus faecium* WHE 81. *International Journal of Food Microbiology* 70:291-301.
- FAO/WHO (2014). Report of a Joint FAO/WHO Working Group on Drafting Guidelines for the evaluation of probiotics in food, Guidelines for the evaluation of probiotics in food. Retrieved from http://www.who.int/foodsafety/fs_management/en/probiotic_guidelines.
- Fernández, M. F., Boris, S. and Barbés, C. (2003). Probiotic properties of human lactobacilli strains to be used in the gastrointestinal tract. *Journal of Applied Microbiology* 94:449-455.
- Ferraro, M. J. (2001). Performance standards for antimicrobial susceptibility testing. USA: NCCLS.
- Flynn, S., Sinderen, D., Thornton, G. M., Holo, H., Nes, I. F. and Collins, J. K. (2002). Characterization of the genetic locus responsible for the production of ABP-118, a novel bacteriocin produced by the probiotic bacterium *Lactobacillus salivarius* subsp. *salivarius* UCC118. *Microbiology* 148:973-984.
- Gómez, N. C., Ramiro, J. M. P., Quecan, B. X. V. and de Melo Franco, B. D. G. (2016). Use of potential probiotic lactic acid bacteria (LAB) biofilms for the control of *Listeria monocytogenes*, *Salmonella Typhimurium*, and *Escherichia coli* O157:H7 biofilms formation. *Frontiers in Microbiology* 7:863.
- Hoque, M., Akter, F., Hossain, K., Rahman, M., Billah, M. and Islam, K. (2010). Isolation, identification and analysis of probiotic properties of *Lactobacillus* spp. from selective regional yoghurts. *World Journal of Dairy and Food Sciences* 5:39-46.

- Ibourahema, C., Dauphin, R. D., Jacqueline, D. and Thonart, P. (2008). Characterization of lactic acid bacteria isolated from poultry farms in Senegal. *African Journal of Biotechnology* 7:2006-2012.
- Jung, J. Y., Lee, S. H., Lee, H. J., Seo, H. Y., Park, W. S. and Jeon, C. O. (2012). Effects of *Leuconostoc mesenteroides* starter cultures on microbial communities and metabolites during kimchi fermentation. *International Journal of Food Microbiology* 153:378-387.
- Kergourlay, G., Messaoudi, S., Dousset, X. and Prévost, H. (2012). Genome sequence of *Lactobacillus salivarius* SMXD51, a potential probiotic strain isolated from chicken cecum, showing anti-campylobacter activity. *Journal of Bacteriology* 194:3008-3009.
- Kingcha, Y., Tosukhowong, A., Zendo, T., Roytrakul, S., Luxananil, P., Chareonpornsook, K., Valyasevi, R., Sonomoto, K. and Visessanguan, W. (2012). Anti-*listeria* activity of *Pediococcus pentosaceus* BCC 3772 and application as starter culture for Nham, a traditional fermented pork sausage. *Food Control* 25:190-196.
- Kirtzalidou, E., Pramateftaki, P., Kotsou, M. and Kyriacou, A. (2011). Screening for lactobacilli with probiotic properties in the infant gut microbiota. *Anaerobe* 17:440-443.
- Kizerwetter-Świda, M. and Binek, M. (2016). Assessment of potentially probiotic properties of *Lactobacillus* strains isolated from chickens. *Polish journal of veterinary sciences* 19:15-20.
- Klayraung, S., Viernstein, H., Sirithunyalug, J. and Okonogi, S. (2008). Probiotic properties of lactobacilli isolated from Thai traditional food. *Scientia Pharmaceutica* 76:485-504.
- Kongkiattikajorn, J. (2015). Potential of starter culture to reduce biogenic amines accumulation in som-fug, a Thai traditional fermented fish sausage. *Journal of Ethnic Foods* 2:186-194.
- Koninkx, J. F. J. G., Tooten, P. C. J. and Malago, J. J. (2010). Probiotic bacteria induced improvement of the mucosal integrity of enterocyte-like Caco-2 cells after exposure to *Salmonella* Enteritidis 857. *Journal of Functional Foods* 2:225-234.
- Langa, S., Maldonado-Barragán, A., Delgado, S., Martín, R., Martín, V., Jiménez, E., Ruíz-Barba, J. L., Mayo, B., Connor, R. I., Suárez, J. E. and Rodríguez, J. M. (2012). Characterization of *Lactobacillus salivarius* CECT 5713, a strain isolated from human milk: from genotype to phenotype. *Applied Microbiology and Biotechnology* 94:1279-1287.
- Liu, S. Q., Asmundson, R., Gopal, P., Holland, R. and Crow, V. (1998). Influence of reduced water activity on lactose metabolism by *Lactococcus lactis* subsp. *cremoris* at different pH Values. *Applied and Environmental Microbiology* 64:2111-2116.
- Liu, Y., Xie, X. X., Ibrahim, S. A., Khaskheli, S. G., Yang, H., Wang, Y. F. and Huang, W. (2016). Characterization of *Lactobacillus pentosus* as a starter culture for the fermentation of edible oyster mushrooms (*Pleurotus* spp.). *LWT - Food Science and Technology* 68:21-26.
- Martín, R., Jiménez, E., Olivares, M., Marín, M. L., Fernández, L., Xaus, J. and Rodríguez, J. M. (2006). *Lactobacillus salivarius* CECT 5713, a potential probiotic strain isolated from infant feces and breast milk of a mother-child pair. *International Journal of Food Microbiology* 112:35-43.
- Menconi, A., Kallapura, G., Latorre, J. D., Morgan, M. J., Pumford, N. R., Hargis, B. M. and Tellez, G. (2014). Identification and characterization of lactic acid bacteria in a commercial probiotic culture. *Bioscience of Microbiota, Food and Health* 33:25-30.

- Messaoudi, S., Manai, M., Kergourlay, G., Prevost, H., Connil, N., Chobert, J. M. and Dousset, X. (2013). *Lactobacillus salivarius*: bacteriocin and probiotic activity. *Food Microbiology* 36:296-304.
- Narakaew, T., Pilasombut, K., Ngamyeesoon, N. and Swetwiwathana, A. (2010). Preliminary characterization of *Lactobacillus salivarius* K7 for probiotic properties. *KKU Research Journal* 15:878-888.
- Neville, B. and O'Toole, P. (2010). Probiotic properties of *Lactobacillus salivarius* and closely related *Lactobacillus* species. *Future Microbiology* 5:759-774.
- Nouri, M., Rahbarizadeh, F., Ahmadvand, D., Moosakhani, F., Sadeqzadeh, E., Lavasani, S. and Khoddami Vishteh, V. (2010). Inhibitory effects of *Lactobacillus salivarius* and *Lactobacillus crispatus* isolated from chicken gastrointestinal tract on *Salmonella* Enteritidis and *Escherichia coli* growth. *Iranian Journal of Biotechnology* 8:32-37.
- O'Shea, E. F., O'Connor, P. M., Raftis, E. J., O'Toole, P. W., Stanton, C., Cotter, P. D., Ross, R. P. and Hill, C. (2011). Production of multiple bacteriocins from a single locus by gastrointestinal strains of *Lactobacillus salivarius*. *Journal of Bacteriology* 193:6973-6982.
- Pilasombut, K., Sakpuaram, T., Wajjwalku, W., Nitisinprasert, S., Swetwiwathana, A., Zendo, T., Fujita, K., Nakayama, J. and Sonomoto, K. (2006). Purification and amino acid sequence of bacteriocins produced by *Lactobacillus salivarius* K7 isolated from chicken intestine. *Songklanakarin Journal of Science and Technology* 28:121-131.
- Ranganath, S. and Sharmila, T. (2016). Isolation and identification of a bacteriocin-producing bacteria from various sources. *International Journal of Current Microbiology and Applied Sciences* 5:655-664.
- Rumjuankiat, K., Keawsompong, S. and Nitisinprasert, S. (2017). Bacterial contaminants from frozen puff pastry production process and their growth inhibition by antimicrobial substances from lactic acid bacteria. *Food Science and Nutrition* 5:454-465.
- Sahadeva, R., Leong, S., Chua, K., Tan, C., Chan, H., Tong, E., Wong, S. and Chan, H. (2011). Survival of commercial probiotic strains to pH and bile. *International Food Research Journal* 18:1515-1522.
- Saint-Cyr, M. J., Haddad, N., Taminiou, B., Poezevara, T., Quesne, S., Amelot, M., Daube, G., Chemaly, M., Dousset, X. and Guyard-Nicodème, M. (2017). Use of the potential probiotic strain *Lactobacillus salivarius* SMXD51 to control *Campylobacter jejuni* in broilers. *International Journal of Food Microbiology* 247:9-17.
- Saithong, P., Panthavee, W., Boonyaratanakornkit, M. and Sikkhamondhol, C. (2010). Use of a starter culture of lactic acid bacteria in plaasom, a Thai fermented fish. *Journal of Bioscience and Bioengineering* 110:553-557.
- Sanders, M. E., Akkermans, L. M. A., Haller, D., Hammerman, C., Heimbach, J. T., Hörmannspurger, G. and Huys, G. (2010). Safety assessment of probiotics for human use. *Gut Microbes* 1:164-185.
- Shehata, M. G., El Sohaimy, S. A., El-Sahn, M. A. and Youssef, M. M. (2016). Screening of isolated potential probiotic lactic acid bacteria for cholesterol lowering property and bile salt hydrolase activity. *Annals of Agricultural Sciences* 61:65-75.
- Swetwiwathana, A. and Visessanguan, W. (2015). Potential of bacteriocin-producing lactic acid bacteria for safety improvements of traditional Thai fermented meat and human health. *Meat Science* 109:101-105.
- Therdthatha, P., Tandumrongpong, C., Pilasombut, K., Matsusaki, H., Keawsompong, S. and Nitisinprasert, S. (2016). Characterization of antimicrobial substance from *Lactobacillus salivarius* KL-D4 and its application as biopreservative for creamy filling. *SpringerPlus* 5:1060.

- Tinrat, S., Saraya, S. and Chomnawang, M. T. (2011). Isolation and characterization of *Lactobacillus salivarius* MTC 1026 as a potential probiotic. *The Journal of general and applied microbiology* 57:365-378.
- Tulumoglu, S., Yuksekdog, Z. N., Beyatli, Y., Simsek, O., Cinar, B. and Yaşar, E. (2013). Probiotic properties of lactobacilli species isolated from children's feces. *Anaerobe* 24:36-42.
- Walsh, M. C., Gardiner, G. E., Hart, O. M., Lawlor, P. G., Daly, M., Lynch, B., Richert, B. T., Radcliffe, S., Giblin, L. and Hill, C. (2008). Predominance of a bacteriocin-producing *Lactobacillus salivarius* component of a five-strain probiotic in the porcine ileum and effects on host immune phenotype. *FEMS microbiology ecology* 64:317-327.
- Wang, R., Jiang, L., Zhang, M., Zhao, L., Hao, Y., Guo, H., Sang, Y., Zhang, H. and Ren, F. (2017). The adhesion of *Lactobacillus salivarius* REN to a human intestinal epithelial cell line requires S-layer proteins. *Scientific Reports* 7:44029.
- Zárate, G., Chaia, A. P., González, S. and Oliver, G. (2000). Viability and β -galactosidase activity of dairy propionibacteria subjected to digestion by artificial gastric and intestinal fluids. *Journal of Food Protection* 63:1214-1221.
- Zhao, D., Du, R. P., Ge, J. P., Wang, X. Y., Mu, L. Q. and Ping, W. X. (2016). Impact of *Lactobacillus paracasei* HD1.7 as a starter culture on characteristics of fermented Chinese cabbage (*Brassica rapa* var. *pekinensis*). *Food Science and Technology Research* 22:325-330.

(Received: 19 September 2017, accepted: 30 October 2017)