

---

## Differential gene expression during flower bud initiation and flower bud development of *Rhynchostylis gigantea* (Lindl.) Ridl

---

**Bounma Phengphachanh<sup>\*</sup>, Daruni Naphrom, Weenun Bundithya and Nuttha Potapohn**

Department of Plant Science and Natural Resources, Faculty of Agriculture, Chiang Mai University, Chiang Mai 50200, Thailand.

Bounma Phengphachanh, Daruni Naphrom, Weenun Bundithya and Nuttha Potapohn (2013). Differential gene expression during flower bud initiation and flower bud development of *Rhynchostylis gigantea* (Lindl.) Ridl. International Journal of Agricultural Technology 9(5):1285-1295.

**Abstract** Generally, *Rhynchostylis gigantea* (Lindl.) Ridl usually induce flowers during winter time, January-February. There was a report which stated that growing *R. gigantea* under short day condition could induce flowering faster than natural flowering period. Thus, this study was conducted to determine the gene expression during floral induction of this plant. Plants were grown under natural condition and short day condition. Shoot tips of those plants were collected every five days, then, each sample was used for total RNA extraction and followed with cDNA synthesis. Fifty- eight arbitrary primers with oligo dT<sub>12</sub>VA, dT<sub>12</sub>VC, dT<sub>12</sub>VG and dT<sub>12</sub>VT were screened. It was found that combinations of 2 primers with dT<sub>12</sub>VC, 8 primers with dT<sub>12</sub>VG and 12 primers with dT<sub>12</sub>VT were able to amplify polymorphic bands differentially expressed among treatments. Selected bands were cloned and sequenced. Thirty TDFs from these combinations, dT<sub>12</sub>VG+OPF16; dT<sub>12</sub>VC+OPD07; dT<sub>12</sub>VT+OPAB17 and dT<sub>12</sub>VT+OPF13 were retrieved for reamplification and cloning. Only 19 TDFs were subsequently sequenced. The sequence analyses revealed that TDFB1, TDFC5, TDFC6 and TDFC8 shared the homology with different species of *Pseudomonas*. TDFC5 and TDFC8 shared the homology with dairy cattle and marine animal, whereas 9 TDFBs (TDFB2 to TDFB10) shared the homology with 1) flower-specific gene promoters of *Oncidium Gower Ramsey*, 2) the transcripts encoding protein gene of *Phalaenopsis equestris* and 3) the vernalization-induced floral transition gene of *Dendrobium nobile*. The other 4 TDFs of TDFD shared the homology with mixed tissue cDNA library of *Nicotiana benthamiana*. On the other hand, TDFC1 and TDFC3 had no significant sequence homology in the database.

This study could suggest that short day was a favorable factor affecting flower bud initiation. During that period, TDFB2 to TDFB10 were detected which could be related to flower bud initiation of *R. gigantea*.

**Keywords:** *Rhynchostylis gigantea* (Lindl.) Ridl, day length, Gene expression, flowering.

---

<sup>\*</sup> **Corresponding author:** Bounma Phengphachanh; e-mail: [bm\\_pheng@hotmail.com](mailto:bm_pheng@hotmail.com)

## Introduction

Generally, plants can flower when they have reached a certain stage of growth or received proper conditions, e.g., temperature and photoperiod, that can induce flowering process. The initiation of flowering is the first step in flower development which can be triggered by environmental and endogenous factors (Taiz and Zeiger, 2007; Amasino and Michaels, 2010). There has been quite a number of information on flower induction of orchid species, e.g., in vitro flowering of several *Dendrobium* hybrids (Goh, 1996); thermoperiodism on flowering of *Dendrobium* Second Love (Campos and Kerbauy, 2004); cool temperature on flowering of most of *Phalaenopsis* hybrids (Lopez *et al.*, 2007) and day length and temperature as well as GA<sub>3</sub> application on flowering of *Rhynchostylis gigantea* (Theesoda, 2004; Talee, 2008; Banyai, 2009; Watthanasrisong, 2010; Phengphachanh *et al.*, 2012).

The transition from vegetative to reproductive growth is controlled by genes. The flower initiation of *Arabidopsis thaliana* occurs in the leaves and induces systemic signals. The signalling is activated by floral promoter gene, CONSTANS (CO) and directly activates the expression of transcription factor gene, FLOWER LOCUS T (FT). FT-protein is synthesized and translocated to shoot apical meristem (SAM) to initiate flowering. FLOWER LOCUS D (FD) is formed at SAM and flowering occurs by repressing floral identity genes, APETALA 1 (AP1) and SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1 (SOC 1) (Amasino and Michaels, 2010).

Several genes which control flowering in orchids have been identified. In *Dendrobium* Madame Thong-In under in vitro condition, DOMADS 1, DOMADS 2 and DOMADS 3 were activated during floral transition (Yu and Goh, 2000). Yu *et al.* (2002) reported that DOMADS1 was a marker gene and specifically expressed in the transitional shoot apical meristem during floral transition in *D. Madame Thong-In*. For *Oncidium* Gower Ramsey, OnTI1, OnTI2 and OnTI3 were highly expressed at different stages of flower development (Hsu *et al.*, 2011). In *Dendrobium nobile*, DnVRN1 gene, which was related to the induction of flowering, was detected in vernalized axillary buds at the middle stage of vernalization (Liang *et al.*, 2012).

As a part of the study on flowering, this work was conducted in order to determine genes which are differentially expressed during floral transition process and flower development of *Rhynchostylis gigantea*.

## **Materials and methods**

### ***Plant materials***

*Rhynchostylis gigantea* (Lindl.) Ridl., 2-3 years old plants, which flowered in the previous year, were grown under natural condition and short day condition (10 h of light) for 3 months. After that, all plants were transferred to the shaded house for further growth and development. The experiment was conducted during May 2<sup>nd</sup>, 2011 to February 2<sup>nd</sup>, 2012, at Orchid Nursery, Mae Hia Agricultural Center for Research, Demonstration and Training, Department of Plant Science and Natural Resources, Faculty of Agriculture, Chiang Mai University, Thailand.

According to the experiment of Talee (2008), the first visible flower bud appeared around 30 days after *R. gigantea* plants were grown under short day regardless of GA<sub>3</sub> application at different concentrations, 1000 and 3000 ppm. Thus, in this study, shoot samples were randomly taken on day-0 and then every 5 days until day 35<sup>th</sup>. Three plants per treatment from each of the 2 treatments, natural and short day condition were used. After that, samples were collected every 20 days until day 95<sup>th</sup>.

### ***Identification of gene expression***

#### ***RNA Extraction***

Total RNA was extracted from fresh shoot tip, using TRIzol<sup>®</sup> Reagent (Invitrogen<sup>™</sup>) following the manufacturer's instructions. The RNA pellet was dissolved in 40 µl of Ultra Pure<sup>™</sup> distilled water prior to DNase treatment. DNase-free RNA (5µl) were reverse transcribed for first strand cDNA, using SuperScript<sup>®</sup> III First-Strand Synthesis System for RT-PCR (Invitrogen<sup>™</sup>) with some modifications. Single strand cDNA was separated into 4 groups which could be tagged with 4 primers, i.e., oligo dT<sub>12</sub>VA, dT<sub>12</sub>VC, dT<sub>12</sub>VG and dT<sub>12</sub>VT.

#### ***Primers screening***

Fifty-eight random primers, i.e., OPD 01- OPD 20; OPF 01- OPF 20; AP-A 01- AP-A10; OPAB 11- OPAB 14 and OPAB 16- OPAB 19, were screened with each of the 4 tagged cDNA groups. Two µl (100ng/µl) of cDNA templates were mixed with reaction buffer according to Anuchai (2011) and Chaipar (2011) procedures. Polymerase Chain Reaction (PCR) was performed, started with 3 mins for activation at 94 °C, followed by 40 cycles of 94 °C for 30 sec,

42 °C for 1 min, and 72 °C for 1 min and extended at 72 °C for 8 mins. The amplified product was electrophoresed on 6% polyacryamide gel at 200 mA, 1200 V and 50 W for 5 h. The gel was silver-stained and dried under the vacuum at 60 °C for 3 h. Primers which yielded polymorphism were selected for further steps.

### ***Re-amplification and cloning***

Potential bands amplified by selected primers were excised and eluted in 20 µl of TE buffer at room temperature for 10 mins, ground using micro-tip and incubated for 15 mins at 95 °C using heat boxes and kept at -20 °C overnight. The re-amplification was done using 2 µl of eluted template under the same reaction and PCR conditions previously mentioned. The re-amplified products were electrophoresed on 1.2% agarose gel at 100 V for 40 mins. The gel was stained using 1% of ethidium bromide and photographed under ultra violet light (Gel Documentations, Lab Focus). The bands of expected size were selected and subsequently cloned into pGEM<sup>®</sup>-T Easy Vector Systems II (Promega) following the manufacturer's instructions. Plasmids containing expected inserts were submitted to Pacific Science Co., LTD (<http://www.pacific-science.co.th>) for sequencing. All sequences were analyzed and compared with all known database from GenBank, using BLAST and ETS programs (<http://www.ncbi.nlm.nih.gov>).

### **Results**

*R. gigantea* plants were grown under natural condition and short day condition for 3 months, after that all plants were transferred to natural condition, under shaded house. Under short day condition, 70% of plants could give visible flower buds at approximately 140 days after treatment whereas 79% of plants under natural condition could give visible flower buds at approximately 163 days after treatment. Flower bud development of all plants grown under natural condition and short day condition could yield flowers and all flowers senesced around the same time.

### ***Screening of primers***

Fifty-eight primers were used for primer screening. The results revealed that twenty primers could providing polymorphic bands. Eight primers, OPD10, OPD18, OPD20, OPF11, A-PA07, OPAB13, OPAB16 and OPAB17 yielded polymorphism with oligo dT<sub>12</sub>VG-tagged samples. Two primers, OPD07 and OPF16 yielded polymorphism with oligo dT<sub>12</sub>VC-tagged samples.

Twelve primers, OPD05, OPD16, OPF04, OPF06, OPF09, OPF12, OPF13, OPF14, OPAB12, OPAB16, OPAB17 and OPAB18 yielded polymorphism with oligo dT<sub>12</sub>VT-tagged samples. OPAB16 and OPAB17 primers yielded polymorphism with oligo dT<sub>12</sub>VG and oligo dT<sub>12</sub>VT-tagged samples. Samples which were tagged with oligo dT<sub>12</sub>VA showed barely or weak band. These 20 primers were used to determine the differential gene expression during flower bud initiation and flower bud development of *R. gigantea*. Result showed that the combinations of oligo dT<sub>12</sub>VG+OPF16, oligo dT<sub>12</sub>VC+OPD07 and oligo dT<sub>12</sub>VT+OPAB17 could yield polymorphic bands during flower bud initiation stage whereas oligo dT<sub>12</sub>VT+OPF13 could yield polymorphic bands during flower bud development stage.

The combination of oligo dT<sub>12</sub>VG+OPF16 could yield bands at the range of 204 - 2,520 bp. The bands were visible for almost all samples. However, the 204 bp band was specific for short day condition on day 5<sup>th</sup>, 10<sup>th</sup> and 35<sup>th</sup> after treatment. These 3 bands were selected for further re-amplification and cloning (Table 1).

The combination of oligo dT<sub>12</sub>VC+OPD07 could yield bands at the range of 1,534 - 3,439 bp. The bands were visible for almost all samples. However, the 1,534 bp band was specific for natural condition on day 5<sup>th</sup>, 10<sup>th</sup>, 20<sup>th</sup>, 25<sup>th</sup>, 30<sup>th</sup> and 35<sup>th</sup> and for short day condition on day 10<sup>th</sup>, 15<sup>th</sup>, 30<sup>th</sup> and 35<sup>th</sup> after treatment. These 10 bands were selected for further re-amplification and cloning (Table 1).

The combination of oligo dT<sub>12</sub>VT+OPAB17 could yield bands at the range of 367- 658 bp. The bands were visible for almost all samples. The 476 and 428 bp bands were specific for natural condition on day 25<sup>th</sup> after treatment and for short day condition on day 5<sup>th</sup>, 10<sup>th</sup>, 20<sup>th</sup> and 25<sup>th</sup> after treatment. These 9 bands were selected for further re-amplification and cloning (Table 1).

The combination of oligo dT<sub>12</sub>VT+OPF13 could yield bands at the range of 4,081-5,847 bp. The bands were visible for almost all samples. The 4,081 and 5,115 bp bands were specific for natural condition on day 55<sup>th</sup>, 75<sup>th</sup> and 95<sup>th</sup> after treatment and for short day condition on day 55<sup>th</sup> after treatment. These 8 bands were selected for further re-amplification and cloning (Table 1).

### ***Analysis of sequences***

Thirty transcript-derived fragments (TDF) were reamplified, and 19 TDFs, 10 of TDFB, 4 of TDFC and 5 of TDFD, showed expected sizes. These 19 TDFs were sequenced (Table 2). TDFB1 and TDFC6 were matched to *Pseudomonas fluorescens* with 82-91% homology. TDFB2 to TDFB10 were matched to orchids genes with 93-94 % homology.

TDFC1 and TDFC3 had no significant sequence homology in the database. TDFC5 and TDFC8 were matched to dairy cattle genes with 94% homology and marine animal *Hydractinia echinata* with 95% homology. TDFD2 to TDFD8 were matched with *Nicotiana benthamiana* sequence with 93-96% homology.

**Table 1.** Primer combination, treatment and ID of selected clones

| Combination                    | Treatment                 | Clone ID  | Combination                    | Treatment | Clone ID  |       |
|--------------------------------|---------------------------|-----------|--------------------------------|-----------|-----------|-------|
| oligo dT <sub>12</sub> VG<br>+ | SD at D5                  | TDFA1     | Oligo dT <sub>12</sub> VT<br>+ | SD at D10 | TDFC3     |       |
|                                | SD at D10                 | TDFA2     |                                | SD at D10 | TDFC4     |       |
|                                | SD at D35                 | TDFA3     |                                | SD at D20 | TDFC5     |       |
| OPF16                          | NC at D5                  | TDFB1     | OPAB17                         | NC at D25 | TDFC6     |       |
|                                | NC at D10                 | TDFB2     |                                | NC at D25 | TDFC7     |       |
|                                | SD at D10                 | TDFB3     |                                | SD at D25 | TDFC8     |       |
|                                | SD at D15                 | TDFB4     |                                | SD at D25 | TDFC9     |       |
|                                | Oligo dT <sub>12</sub> VC | NC at D20 |                                | TDFB5     | NC at D55 | TDFD1 |
|                                | +                         | NC at D25 |                                | TDFB6     | NC at D55 | TDFD2 |
|                                | OPD07                     | NC at D30 |                                | TDFB7     | SD at D55 | TDFD3 |
| Oligo dT <sub>12</sub> VT<br>+ | SD at D30                 | TDFB8     | Oligo dT <sub>12</sub> VT<br>+ | SD at D55 | TDFD4     |       |
|                                | NC at D35                 | TDFB9     |                                | NC at D75 | TDFD5     |       |
|                                | SD at D35                 | TDFB10    |                                | NC at D75 | TDFD6     |       |
|                                | OPF13                     |           |                                | NC at D95 | TDFD7     |       |
| Oligo dT <sub>12</sub> VT      | SD at D5                  | TDFC1     |                                | NC at D95 | TDFD8     |       |
| +                              | SD at D5                  | TDFC2     |                                |           |           |       |
| OPAB17                         |                           |           |                                |           |           |       |

## Discussion

*Rhynchostylis gigantea* plants were grown under different conditions, short day and natural conditions. It was found that plants grown under short day could give flower 23 days earlier than those grown under natural condition. That means flower initiation should have occurred at different time, resulting from earlier induction by short day condition.

Sequence analysis of differentially-expressed TDFs revealed positive matches to the database. Nine TDFs, i.e., TDFB2, TDFB3, TDFB4, TDFB5, TDFB6, TDFB7, TDFB8, TDFB9 and TDFB10, were matched with orchid genes, i.e., 1) flower-specific gene promoters of *Oncidium Gower Ramsey*, from flower organ (Hsu *et al.*, 2011), 2) the transcripts encoding protein gene of *Phalaenopsis equestris*, gene from flower bud (Tsai *et al.*, 2006) and 3) the vernalization-induced floral transition gene of *Dendrobium nobile* (20+30) d-vernalized cDNA library, detecting from axillary buds of *D. nobile* at middle stage of vernalization (Liang *et al.*, 2012). These TDFs were found in plants grown under short day and natural conditions with some differences. Under short day, TDFs could be detected

on days 10<sup>th</sup>, 15<sup>th</sup>, 30<sup>th</sup> and 35<sup>th</sup> whereas TDFs of natural condition could be found on days 10<sup>th</sup>, 20<sup>th</sup>, 25<sup>th</sup>, 30<sup>th</sup> and 35<sup>th</sup>. In addition, TDFA and TDFC could be noticed only in plants grown under short day condition. Unfortunately, ones from TDFA could not be reamplified and cloned due to technical difficulties. These TDFAs are very interesting in that the expression profiles could be involved with flowering since they were found in plants grown under short day at days 5<sup>th</sup>, 10<sup>th</sup> and 15<sup>th</sup>, but none was present in plant grown under natural condition. Further investigation regarding this type of TDF should be conducted to elucidate gene expression controlling flower induction and initiation.

Other TDF results showed general information on living things. The results matched to dairy cattle and marine animal. However, the result from TDFB1 and TDFC6 showed matching with the complete genome of *Pseudomonas fluorescens* Pf-5 and SBW25, respectively. *P. fluorescens* are common soil bacteria that can improve plant health through nutrient cycling, pathogen antagonism and induction of plant defenses by producing secondary metabolites that suppress plant pathogens (Paulsen *et al.*, 2005; Silby *et al.*, 2009). Interestingly, *R. gigantea* is an epiphytic orchid and gene expression from this species is similar to those found in plants grown in soil. This might help explain why this type of orchid can survive without any growing medium.

**Table 2.** Expression pattern, reamplification and sequence search of selected clones

| Day after treatment | TDFA  |           | TDFB  |           | TDFC   |           | TDFD  |           |
|---------------------|---|-----------|---|-----------|--|-----------|---|-----------|
|                     | oligodT <sub>12</sub> VG+OPF16<br>Natural condition | Short day | oligodT <sub>12</sub> VC+OPD07<br>Natural condition | Short day | oligodT <sub>12</sub> VT+OPAB17<br>Natural condition | Short day | oligodT <sub>12</sub> VT+OPF13<br>Natural condition | Short day |
| D0                  |   |           |   |           |  |           |   |           |
| D5                  | Δ   | √         |   |           |  | O ⊗       |   |           |
| D10                 | Δ   | √         |   | √         |  | O ⊗       |   |           |
| D15                 |   |           |   | √         |  |           |   |           |
| D20                 |   |           | √   |           |  | √         |   |           |
| D25                 |   |           | √   |           | √ ⊗  | √ ⊗       |   |           |
| D30                 |   |           | √   | √         |  |           |   |           |
| D35                 | Δ   | √         | √   | √         |  |           |   |           |
| D55                 |   |           |   |           |  |           | ⊗ √   | ⊗ √       |
| D75                 |   |           |   |           |  |           | ⊗ √   |           |
| D95                 |   |           |   |           |  |           | ⊗ √   |           |

Note: Δ: could not re-amplify; √: matched to database; O: not match to any information in data base; ⊗: Not expected size

**Table 3.** Characterization of genes differentially expressed in flower bud initiation and flower buds development of *R. gigantea* (Lindl.) Ridl

| Clone | Sequence size (bp) | Data from gene bank |  |                   |              |
|-------|--------------------|---------------------|--|-------------------|--------------|
|       |                    | Accession Number    | Homology to  | Comparisons score | Homology (%) |
| TDFB1 | 270                | CP000076.1          | Pseudomonas fluorescens Pf-5, complete genome                            | 254               | 84           |
|       |                    | CP002585.1          | Pseudomonas brassicacearum subsp. brassicacearum NFM421, complete genome | 210               | 82           |
|       |                    | CP000075            | Pseudomonas syringae pv. syringae B728a, complete genome                 | 130               | 88           |
| TDFB2 | 271                | HS522463            | Oncidium Gower Ramsey, flower cDNA library                               | 368               | 94           |
|       |                    | CK857414            | Flower bud of Phalaenopsis equestris, cDNA 5', mRNA sequence             | 248               | 94           |
|       |                    | HO199381.1          | Dendrobium nobile (20+30)d-vernalized cDNA library                       | 381               | 94           |
| TDFB3 | 271                | HS522463            | Oncidium Gower Ramsey, flower cDNA library                               | 368               | 94           |
|       |                    | CK857414            | Flower bud of Phalaenopsis equestris, cDNA 5', mRNA sequence             | 248               | 94           |
|       |                    | HO199381.1          | Dendrobium nobile (20+30)d-vernalized cDNA library                       | 381               | 94           |
| TDFB4 | 271                | HS522463            | Oncidium Gower Ramsey, flower cDNA library                               | 368               | 94           |
|       |                    | CK857414            | Flower bud of Phalaenopsis equestris, cDNA 5', mRNA sequence             | 248               | 94           |
|       |                    | HO199381.1          | Dendrobium nobile (20+30)d-vernalized cDNA library                       | 381               | 94           |
| TDFB5 | 271                | HS522463            | Oncidium Gower Ramsey, flower cDNA library                               | 368               | 94           |
|       |                    | CK857414            | Flower bud of Phalaenopsis equestris, cDNA 5', mRNA sequence             | 248               | 94           |
|       |                    | HO199381.1          | Dendrobium nobile (20+30)d-vernalized cDNA library                       | 381               | 94           |
| TDFB6 | 272                | HS522463            | Oncidium Gower Ramsey, flower cDNA library                               | 374               | 93           |
|       |                    | CK857414            | Flower bud of Phalaenopsis equestris, cDNA 5', mRNA sequence             | 248               | 94           |



|        |     |            |  |      |    |
|--------|-----|------------|--|------|----|
|        |     | HO199381.1 | Dendrobium nobile<br>(20+30)d-vernalized cDNA library        | 387  | 94 |
| TDFB7  | 271 | HS522463   | Oncidium Gower Ramsey, flower cDNA library                   | 363  | 93 |
|        |     | CK857414   | Flower bud of Phalaenopsis equestris, cDNA 5', mRNA sequence | 243  | 93 |
|        |     | HO199381.1 | Dendrobium nobile<br>(20+30)d-vernalized cDNA library        | 375  | 94 |
| TDFB8  | 271 | HS522463   | Oncidium Gower Ramsey, flower cDNA library                   | 363  | 93 |
|        |     | CK857414   | Flower bud of Phalaenopsis equestris, cDNA 5', mRNA sequence | 248  | 94 |
|        |     | HO199381.1 | Dendrobium nobile<br>(20+30)d-vernalized cDNA library        | 375  | 93 |
| TDFB9  | 271 | HS522463   | Oncidium Gower Ramsey, flower cDNA library                   | 368  | 94 |
|        |     | CK857414   | Flower bud of Phalaenopsis equestris, cDNA 5', mRNA sequence | 248  | 94 |
|        |     | HO199381.1 | Dendrobium nobile<br>(20+30)d-vernalized cDNA                | 381  | 94 |
| TDFB10 | 271 | HS522463   | Oncidium Gower Ramsey, flower cDNA library                   | 368  | 94 |
|        |     | CK857414   | Flower bud of Phalaenopsis equestris, cDNA 5', mRNA sequence | 248  | 94 |
|        |     | HO199381.1 | Dendrobium nobile<br>(20+30)d-vernalized cDNA                | 381  | 94 |
| TDFC1  | 476 |            | No significant sequence homology in the database             |      |    |
| TDFC3  | 476 |            |  |      |    |
| TDFC5  | 476 | AV725203   | Bos taurus cDNA clone  | 472  | 94 |
|        |     | DN602668   | Hypothetical protein   | 95.3 | 95 |
| TDFC6  | 476 | AM181176   | Pseudomonas fluorescens SBW25 complete genome                | 366  | 91 |
|        |     | CP003041   | Pseudomonas fluorescens A506, complete genome                | 322  | 88 |
| TDFC8  | 555 | AV725203   | Bos taurus cDNA clone  | 472  | 94 |
|        |     | DN602668   | Hypothetical protein of Hydractina echinata                  | 93.5 | 95 |
| TDFD2  | 475 | CK285246   | Nicotiana benthamiana  | 263  | 93 |
|        |     | CK285245   | mixed tissue cDNA library                                    | 84.2 | 96 |
| TDFD4  | 475 | CK285246   | Nicotiana benthamiana  | 263  | 93 |
|        |     | CK285245   | mixed tissue cDNA library                                    | 84.2 | 96 |
| TDFD6  | 475 | CK285246   | Nicotiana benthamiana  | 263  | 93 |
|        |     | CK285245   | mixed tissue cDNA library                                    | 84.2 | 96 |
| TDFD8  | 476 | CK285246   | Nicotiana benthamiana  | 263  | 93 |
|        |     | CK285245   | mixed tissue cDNA library                                    | 84.2 | 96 |

## Conclusion

The identification of gene expression during flower bud initiation and flower bud development was studied. Fifty-eight arbitrary primers were screened on cDNA of *R. gigantea*. Twenty primers showed polymorphisms. Combinations of oligo dT<sub>12</sub>VG+OPF16, oligo dT<sub>12</sub>VC+OPD07, oligo dT<sub>12</sub>VT+OPAB17 yielded polymorphic bands at flower bud initiation while oligo dT<sub>12</sub>VT+OPF13 at flower bud development. Thirty TDFs from these 4 combinations were reamplified, cloned and sequenced.

The sequence analysis revealed TDFB1, TDFC5, TDFC6 and TDFC8 having homology with different species of *Pseudomonas*. TDFC5 and TDFC8 had homology with dairy cattle and marine animal. Nine TDFBs (TDFB2 to TDFB10) shared the homology with flower-specific gene promoters of *Oncidium Gower Ramsey*; the transcripts encoding protein gene of *P. equestris* and the vernalization- induced floral transition gene of *D. nobile*. Four TDFDs shared sequences with mixed tissue cDNA library of *N. benthamiana*. TDFC1 and TDFC3 had no significant sequence homology in the database.

This primary information on molecular practice would guide the study on the flowering process of *R. gigantea* in the future. It could facilitate further characterization of the key players in flowering process pathways. Further study is needed in order to understand and determine genes controlling flower induction, initiation and development of *R. gigantea*.

## Acknowledgements

We gratefully acknowledge ACMECS for financial support. Also thanks are due to the Centre of Excellence on Agricultural Biotechnology, Faculty of Agriculture, Chiang Mai University for providing the facility and information, and DEFECO who supported the plant materials.

## References

- Anuchai, J. (2011). Detection of differential gene expression in sepal and petal of *Spathoglottis* species by DDRT-PCR technique. MSc. Thesis in Horticulture, Chiang Mai University, Chiangmai. Thailand.
- Amasino, R.M. and Michaels, S.D. (2010). The timing of flowering in the future perspectives in plant biology. *Plant Physiol.* 154:516-520.
- Banyai, K. (2009). The effects of gibberellins and light on plants nutrients and off-season flowering of *Rhynchostylis gigantea* (Lindl) Ridl. MSc.Thesis, Chiang Mai University, Thailand.
- Campos, K.O. and Kerbauy, G.B. (2004). Thermoperiodic effect on flowering and endogenous hormonal status in *Dendrobium* (Orchidaceae). *J. Plant Physiol.* 161:1385-1387.
- Chaipar, T. (2011). Detection of differential gene expression in bract of *Curcuma petiolata* Roxb. by DDRT-PCR technique. MSc. Thesis in Horticulture, Chiang Mai University, Chiangmai. Thailand.
- Goh, C.J. (1996). Production of flowering orchid seedling and plantlets. *Malay Orchid Rev.* 30:27-30.

- Hsu, C.T., Liao, D.C., Wu, F.H., Liu, N.T., Shen, S.C., Chou, S.J., Tung, S.Y., Yang, C.H., Chan, M.T. and Lin, C.S. (2011). Integration of molecular biology tools for identifying promoters and genes abundantly expressed in flower of *Oncidium Gower Ramsey*. *BMC. Pl. Bio.* 11:1-14.
- Liang, S., Ye, Q.S., Li, R.H., Leng, J.Y. and Li, M.R. (2012). Transcriptional regulations on the low-temperature-induced floral transition in an Orchidaceae species, *Dendrobium nobile*: An expressed sequence tags analysis. Hindawi Publishing Corporation, pp. 1-14.
- Lopez, R., Runkle, E., Wang, Y.T., Blanchard, M. and Hsu. T. (2007). Orchids march 2007. Available in <http://www.aos.org>. revised January 15: 2012, pp. 182.
- Paulsen, I.T., Press, C.M., Ravel, J., Kobayashi, D.Y., Myers, G.S., Mavrodi, D.V., Deboy, R.T., Seshadri, R., Ren, Q., Madupu, R., Dodson, R.J., Dunrkin, A. S., Brinkac. L.M., Daugherty, S.C., Sullivan, S.V., Rosovitz, M.J., Gwinn, M.L., Zhou, L., Schneider, D.J., Cartinhour, S. W., Nelson, W.C., Weidman, J., Watkins, K., Tran, K., Khouri, H., Pierson, E.A., Pierson, L.S., Thomashow, L.S. and Loper, J.E. (2005). Complete genome sequence of the plant commensal *Pseudomonas fluorescens Pf-5*. *Nat. Biotech.* 7:873-878.
- Phengphachanh, B., Naphrom, D., Bundithya, W. and Potapohn, N. (2012). Effects of day length and gibberellic acid (GA<sub>3</sub>) on flowering and endogenous hormone levels in *Rhynchosytilis gigantea* (Lindl.) Ridl. *Agri. Sci.* 4:217-222.
- Silby, M.W., Cerdeño-Tarraga, A.M., Vernikos, G.S., Giddens, S.R., Jackson, R.W., Preston, G.M., Zhang, X.X., Moon, C.D., Gehrig, S.M., Godfrey, S.A., Knight, C.G., Malone, J. G., Robinson, Z., Spiers, A.J., Harris, S., Challis, G.L., Yaxley, A.M., Harris, D., Seeger, K., Murphy, L., Rutter, S., Squares, R., Quail, M.A., Saunders, E., Mavromatis, K., Brettin, T.S., Bentley, S.D., Hothersall, J., Stephens, E., Thomas, C.M., Parkhill, J., Levy, S.B., Rainey, P.B. and Thomson, N. (2009). Genomic and genetic analyses of diversity and plant interaction of *Pseudomonas fluorescens*. *Genome Biol.* 10:1-16.
- Taiz, L. and Zeiger, E. (2007). *Plant Physiology*. Sinauer Associates, Inc., Publishers: Massachusetts.
- Theesoda, V. (2004). Effects of photoperiod and night temperature on flowering of *Rhynchosytilis gigantea*. Special problem of BSc, Maejo University. Thailand.
- Talee, N. (2008). Effect of short day condition and gibberellic acid on off-season flowering of *Rhynchosytilis gigantea* (Lindl) Ridl. MSc. Thesis. Chiang Mai University. Thailand.
- Tsai, W.C., Hsiao, Y.Y., Lee, S.H., Tung, C.W., Wang, D.P., Wang, H.C., Chen, W.H. and Chen, H.H. (2006). Expression analysis of the ETSS derived from the flower buds of *Phalaenopsis equestris*. *J. Plant Sci.* 170:426-432.
- Yu, H. and Goh, C.J. (2000). Differential gene expression during floral transition in an orchid hybrid *Dendrobium Madame Thong-In*. *Plant Cell Report* 19:926-931.
- Yu, H., Yang S.H. and Goh. C.J. (2002). Spatial and temporal expression of the orchid floral homeotic gene *DOMADS1* is mediated by its upstream regulatory region. *Plant Molec. Biol.* 49:225-237.
- Watthanasrisong, J. (2010). Effects of timing of short day with low temperature on off-season flowering of *Rhynchosytilis gigantea* var. *Harrisonianum* Holtt. MSc. Thesis. Chiang Mai University, Thailand.

(Received 5 July 2013; accepted 31 August 2013)